

Fluctuation of planar Brownian loop capturing large area

Alan Hammond, Yuval Peres

February 1, 2008

Abstract. We consider a planar Brownian loop B that is run for a time T and conditioned on the event that its range encloses the unusually high area of πT^2 , with $T \in (0, \infty)$ being large. The conditioned process, denoted by X , was proposed by Senya Shlosman as a model for the fluctuation of a phase boundary. We study the deviation of the range of X from a circle of radius T . This deviation is measured by the inradius $R_{\text{in}}(X)$ and outradius $R_{\text{out}}(X)$, which are the maximal radius of a disk enclosed by the range of X , and the minimal radius of a disk that contains this range. We prove that, in a typical realization of the conditioned measure, each of these quantities differs from T by at most $T^{2/3+\epsilon}$.

1 Introduction

The goal of this paper is to analyse the fluctuations of a planar Brownian loop under the condition that it encircles a large area. Throughout, $B : [0, T] \rightarrow \mathbb{R}^2$ will denote a standard planar Brownian loop, that is, a planar Brownian motion with initial location $B(0) = 0$ that is conditioned on the event that $B(T) = 0$. Allowing $\text{enc}(B)$ to denote the random set of points that lie in the union of all bounded components of $\mathbb{R}^2 \setminus \{B(t) : t \in [0, T]\}$, our conditioning takes the form

$$|\text{enc}(B)| \geq \pi T^2,$$

where $|\cdot|$ denotes two-dimensional Lebesgue measure. Note that by the spatial-temporal scaling satisfied by Brownian motion, the law of the conditioned process is the same as that obtained from sampling a Brownian loop run for a unit of time that is conditioned to enclose an area exceeding πT , and then dilating space by a factor of $\sqrt{T}$. Throughout, we will define the process on the interval $[0, T]$. The conditioned process will be denoted by $X : [0, T] \rightarrow \mathbb{R}^2$. As we discuss in Section 2, a classical variational principle suggests that the range $X[0, T]$ takes a form close to that of a circle of radius T . The principal aim of this paper is to investigate the magnitude




Figure 1: A sketch of the conditioned motion.

of the deviation of the range $X[0, T]$ from such a circle. Our main theorem provides a bound on a quantity that measures this deviation. To be precise, for a planar compact set K , the inradius $R_{\text{in}}(K)$ of K is the maximal radius of a circle lying in K , while the outradius $R_{\text{out}}(K)$ is the minimal radius of any circle in which K is contained. We will write $R_{\text{in}}(B)$ and $R_{\text{out}}(B)$ for $R_{\text{in}}(\text{enc}(B))$ and $R_{\text{out}}(\text{enc}(B))$, adopting the same shorthand for the process X .

Theorem 1 *Let $\epsilon \in (0, 1/6)$. For $c \in (0, \pi^2/32)$ and all $T \geq T_c$ sufficiently high,*

$$\mathbb{P}\left(R_{\text{in}}(X) < T - T^{\frac{2}{3}+\epsilon}\right) \leq \exp\left\{-cT^{\frac{1}{3}+2\epsilon}\right\},$$

For any constant $\hat{c}$ satisfying $\hat{c} \in (0, \pi^2/2^9)$ and all $T \geq T_{\hat{c}}$ sufficiently high,

$$\mathbb{P}\left(R_{\text{out}}(X) > T + T^{\frac{2}{3}+\epsilon}\right) \leq \exp\left\{-\hat{c}T^{\frac{1}{3}+2\epsilon}\right\}.$$

How close is $X[0, T]$ to the boundary of its convex hull? This is a question about the local nature of the deviation of the conditioned process. We will write $\mathcal{L}(\text{conv}K)$ for the length of the longest line segment that lies in the boundary of the convex hull $\text{conv}(K)$ of K . We also define the maximum local roughness $\text{MLR}(K)$ of K to be the maximal distance between a point in K , and the boundary of $\text{conv}(K)$. That is,

$$\text{MLR}(K) := \sup_{k \in K} \inf_{x \in \partial(\text{conv}(K))} d(x, k).$$

We will write $\mathcal{L}(\text{conv}B)$ for $\mathcal{L}(\text{conv}K)$ in the case where K is the range of the process $B : [0, T] \rightarrow \mathbb{R}^2$. A similar convention will apply for the maximum local roughness, and for the conditioned process $X : [0, T] \rightarrow \mathbb{R}^2$.

Senya Shlosman proposed this model to us, presenting some heuristic arguments that its deviation behaviour has much in common with that observed in numerous models of phase boundaries in two-dimensional random systems, more specifically, that exponents describing the typical behaviour of $\mathcal{L}(\text{conv}X)$ and $\text{MLR}(X)$ coincide with those in these other models. We will present some heuristic arguments of our own in favour of this belief at the end of the introduction. Theorem 1 has the following straightforward consequence.

Corollary 1 *The fluctuation of the conditioned process $X : [0, T] \rightarrow \mathbb{R}^2$ satisfies the following bounds. For any $\epsilon \in (0, 1/12)$, $c \in (0, 2^{-21})$ and for all $T \geq T_c$ sufficiently high,*

$$\mathbb{P}\left(\mathcal{L}(X) \geq T^{\frac{5}{6}+\epsilon}\right) \leq \exp\left\{-cT^{\frac{1}{3}+4\epsilon}\right\},$$

Moreover, for $\epsilon \in (0, 1/6)$ and all $T \geq T_c$ sufficiently high,

$$\mathbb{P}\left(\text{arcl}(\partial(\text{conv}X)) > 2\pi(T + T^{\frac{2}{3}+\epsilon})\right) \leq \exp\left\{-\hat{c}T^{\frac{1}{3}+2\epsilon}\right\},$$

where the constant $\hat{c}$ appears in Theorem 1, and where arcl denotes the arclength of a planar set.

An outline of the proof. We now discuss the techniques required to prove Theorem 1, outlining the structure of the paper as we do so. The first part of Section 2 describes more precisely the assertion from the theory of large deviations that the range $X[0, T]$ resembles a circle of radius T . The second part is devoted to developing the tools required to prove Theorem 1, while the proofs are given in Section 3.

We seek to understand the behaviour of the conditioned motion X by considering the polygon P whose vertices are the locations of X at m equally spaced moments of time. We consider the area of this polygon and that of its convex hull, as well as the area trapped between the range of the motion as it traverses the space between two successive vertices of the polygon, and the line segment between this pair of vertices. We are free to choose the value of m as we please, and it is of little surprise given our belief about the true fluctuation in the model that choices close to $T^{1/3}$ are convenient.

The two most significant results in the second part of Section 2 are Lemma 4 and Proposition 2. The former shows that the convex hull of the polygon P is likely to trap an area that is not much less than that captured by the conditioned motion, for values of m slightly less than $T^{1/3}$. The discrepancy is shown to be at most a little more than $T^{4/3}$ with high probability. Proposition 2 provides an estimate on the regularity of the conditioned motion, to the effect that it is unlikely to move too quickly in short periods of time. More precisely, in a time of order $T^{2/3+\epsilon}$, we rarely see the motion cover as much distance as $T^{2/3+2\epsilon}$. To prove each of

these results, the first step is to provide a lower bound on the probability that the Brownian loop $B : [0, T] \rightarrow \mathbb{R}^2$ in fact satisfies the requirement of the conditioning, that it captures an area of πT^2 . This bound is provided in Lemma 1 by estimating the probability that a regular polygon with an order of $T^{1/3}$ vertices and having this area is enclosed by the motion B . We then prove the two results by showing that if $|\text{conv}(P)|$ is less than $|\text{enc}(X)|$ by $T^{4/3+\epsilon}$, or if X does move a distance of $T^{2/3+2\epsilon}$ in some interval of time of order $T^{2/3+\epsilon}$, then certain functionals of collections of normally distributed random variables assume high values. For example, in the proof of Lemma 4, we consider the event that $|\text{conv}(P)|$ is less than $|\text{enc}(X)|$ by at least $T^{4/3+\epsilon}$. In Lemma 3, we show that the discrepancy $\text{enc}(X) \setminus \text{conv}(P)$ is contained in the union over the polygonal edges l of regions that are, roughly speaking, rectangles whose long axis is l and whose width is the orthogonal fluctuation of the motion X during the interval of time in which it traverses the edge l .

These edge lengths and fluctuations are two collections of random variables whose distributions are readily bounded above by some having a normal distribution. We have seen that the sum of the products of edge lengths and corresponding fluctuations is at least $T^{4/3+\epsilon}$ if the event in question occurs. The square of this expression may be bounded above by the product of two random variables having the χ^2 -distribution, by use of the Cauchy-Schwarz inequality. Computations that make use of the formula for the density of this distribution show that the event in question is less probable than that of the conditioning being satisfied, by a comparison with the lower bound given in Lemma 1.

In the proof of Theorem 1, we work with the polygon P , with m being chosen to be of order $T^{1/3}$. The convex hull of P is known typically to trap a high area by Lemma 4. An isoperimetric result, Lemma 8, gives a lower bound on the arclength of a planar convex body in terms of its area and its global deviation, the latter giving rise to an excess over that occurring in the extremal case of a disk. We deduce that the global deviation of the boundary of the convex hull of P is not too high: for otherwise, the sum of the edge-lengths of the polygon being high implies that the χ^2 -distributed sum of their squares is improbably large. This arclength is also forced to be high if there is any vertex p of P that is too distant from the line segment in $\text{conv}(P)$ which the motion is traversing when it reaches p . The fluctuation of the motion between two successive vertices of the polygon has been bounded in Proposition 2. We have obtained enough control on the motion to deduce that its global deviation typically satisfies an upper bound whose order is comparable to that satisfied by the deviation of the approximating polygon, that is, little more than $T^{2/3}$.

We so demonstrate that the motion is likely to be trapped between circles whose radii differ by an order that slightly exceeds that of $T^{2/3}$. The bounds on $\text{MLR}(X)$ and $\mathcal{L}(X)$ given in Corollary 1 are straightforward consequences.

Remark Throughout, any time parameter takes a value in $[0, T]$. From time to time, terms such as $t + T^{2/3}$ appear that may be greater than T . In such cases, we are referring to the value on $[0, T]$ that is the value stated reduced modulo T .

1.1 Comparisons and heuristic arguments

It is believed that a variety of models of phase boundaries in two-dimensional random systems exhibit the same power-law fluctuations, even though their macroscopic profiles differ. An important example of such a model is that of a large finite cluster in the supercritical phase of the site percolation model in $\mathbb{Z}^2$. Choosing a parameter value $p > p_c$, Alexander and Uzun [7] condition on the event that there exists an open dual circuit surrounding the origin and enclosing an area of at least n^2 , for large n . The asymptotic shape of this circuit is the boundary of a compact convex body, known as the Wulff crystal, that minimises a surface tension, c.f. [10]. The fluctuation of the circuit away from this shape may be measured by the maximum local roughness, which means in this case, the maximum distance of a vertex in the circuit from the boundary of its convex hull. In [1], the maximum local roughness is established to be bounded above by a quantity of the order of $n^{2/3}$. The average local roughness (which is roughly speaking the mean distance of a vertex in C from the convex hull) is bounded above by $n^{1/3}$ up to power order, a bound which is believed to be sharp. In [7], the maximum local roughness is shown to satisfy a lower bound that is given by $n^{1/3}$ if we omit logarithmic corrections.

We expect that the exponents describing the typical behaviour of the two measures of fluctuation, $\mathcal{L}$ and MLR, coincide with those anticipated for the percolation problem. That is, we expect that $\mathcal{L}(\text{conv}X)$ behaves as $T^{2/3}$, and $\text{MLR}(X)$ as $T^{1/3}$. To give an argument that supports the claim that $\mathcal{L}(\text{conv}X)$ is typically not much greater than $T^{2/3}$, suppose that we sample the measure X and find a realization $X(\omega)$ where there is a line segment $L = [x_1, x_2]$ in $\partial(\text{conv}(X))$ whose length exceeds $T^{2/3+\epsilon}$. The times t_1 and t_2 at which $X(\omega)$ visits the endpoints x_1 and x_2 presumably satisfy $|t_1 - t_2| > T^{2/3+\epsilon/2}$, since the conditioned motion tends to move at a fairly constant rate. (Indeed, in Proposition 2, we will prove that the process X is unlikely to cover distances as big as L at speeds significantly greater than the average one at which X moves). Choosing two points t and t^* on the interval $[0, T]$ uniformly and independently of other randomness, we may resample the path of X on the interval $[t, t^*]$, replacing $X[t, t^*]$ by a Brownian bridge that moves from $X(t)$ to $X(t^*)$ in time $t^* - t$. The Markov chain on loops that performs this resampling and jumps to the new path provided that it captures the required area of πT^2 , and stays put in the other case, has the law of the conditioned process X as its invariant measure. We see that, for the action of this resampling on $X(\omega)$, if the points t and t^* happen to be




Figure 2: How resampling may trap more area.

picked near to t_1 and t_2 respectively, then the effect of the resampling is to replace the motion of $X(w)$ as it traverses L by a new motion. This motion has a time of at least $T^{2/3+\epsilon/2}$ to traverse a distance of $T^{2/3+\epsilon}$. This new section of path typically fluctuates orthogonally to L by a distance at least of order $T^{1/3+\epsilon/4}$ (this being the square root of the available time). With a probability that is uniformly bounded below in T , this fluctuation occurs for a fixed but high fraction of time in the direction away from the existing convex hull of X . In this case, the resampled motion would seem to capture an area of the plane that exceeds that captured by $X(w)$ by an amount of the order of $T^{2/3+\epsilon} \cdot T^{1/3+\epsilon/4} = T^{1+5\epsilon/4}$ (the left-hand-side here being the product of the length of L and the orthogonal fluctuation of the resampled motion). In this event, the resampling certainly meets the area criterion. Figure 1.1 shows a sketch of the range of a typical realization of X , and a resampling that creates more trapped area by the means just described. It is very believable that the typical order of the excess of area that X captures over what it must capture, $|\text{enc}(X)| - \pi T^2$, is linear in T , and that

$$\mathbb{P}\left(|\text{enc}(X)| - \pi T^2 \geq T^{1+\alpha}\right) \quad (1)$$

decays at a super-polynomial rate, for any given $\alpha > 0$. However, the preceding argument suggests that the resampled motion - whose law is that

of X - has an excess of area of order of $T^{1+5\epsilon/4}$ with a probability that is at least a polynomially decaying multiple of the probability that there exists a line segment in $\partial(\text{conv}(X))$ of length $T^{2/3+\epsilon}$ (the fact that the points t and t^* must be chosen to be near t_1 and t_2 is responsible for the appearance of a polynomial factor here). So, one expects that the probability of such a line segment in a realization of X decays at a super-polynomial rate in T . It remains an interesting problem to derive such an upper bound on the quantity (1), as does that of obtaining lower bounds on the area captured after resampling (an example of the difficulties involved in determining the area captured is the fact that Lemma 3 is not valid if the instance of $\text{conv}(P)$ in its statement is replaced by P).

2 Understanding how the process X fluctuates

2.1 The macroscopic profile of the range of X

The theory of large deviations is of use in deriving the asymptotic shape of the range of the process X . Let $C_0([0, 1], \mathbb{R}^2)$ denote the space of continuous planar-valued functions $f : [0, 1] \rightarrow \mathbb{R}^2$ for which $f(0) = 0$, and let W denote two-dimensional Wiener measure. Writing $f_T(\cdot) = Tf(\cdot/T)$, we have, by Theorem 5.1 of [8],

$$\limsup_{T \rightarrow \infty} \frac{1}{T} \log W \left\{ g : g \Big|_{[0, T]} = f_T, \text{ for some } f \in C \right\} \leq - \inf_{x \in C} I(x) \quad (2)$$

for $C \subseteq C_0([0, 1], \mathbb{R}^2)$ closed, and

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \log W \left\{ g : g \Big|_{[0, T]} = f_T, \text{ for some } f \in O \right\} \geq - \inf_{x \in O} I(x) \quad (3)$$

for $O \subseteq C_0([0, 1], \mathbb{R}^2)$ open.

The *large deviations' rate function* $I : C_0([0, 1], \mathbb{R}^2) \rightarrow [0, \infty]$ is given by

$$I(f) = \frac{1}{2} \int_0^1 \|Df\|^2(t) dt,$$

if $f \in H_1([0, 1], \mathbb{R}^2)$ (that is, if f is absolutely continuous with square integrable derivative Df), with $I(f) = \infty$ otherwise. By applying the contraction principle of large deviations [3, Section 4.2.1] to the mapping of the space of Brownian paths $Z : [0, T] \rightarrow \mathbb{R}^2$ to the space of Brownian loops $B : [0, T] \rightarrow \mathbb{R}^2$ given by $Z(t) = B(t) - \frac{t}{T}B(T)$, we learn that (2) and (3) are valid for the measure dB provided that the space $C_0([0, 1], \mathbb{R}^2)$ is replaced by its subspace $\overline{C}_0$ consisting of functions f for which $f(1) = 0$. In evaluating the area enclosed by a loop, we will use for the present argument the signed area, given by

$$A(f) = \frac{1}{2} \int_0^1 (f_1(t)f_2'(t) - f_1'(t)f_2(t))dt, \quad (4)$$

for any $f \in H_1([0, 1], \mathbb{R}^2)$. Noting that $A(f_T) \geq \pi T^2$ for such f if and only if $A(f) \geq \pi$, we now identify those functions in $\{f \in H_1([0, 1], \mathbb{R}^2) : A(f) \geq \pi\}$ that minimise $I(f)$. Any f in this set is certainly square integrable, and thus, has an L^2 -convergent complex Fourier series

$$f(x) = \sum_{n \in \mathbb{Z}} a_n \exp\{2n\pi i x\}. \quad (5)$$

Note that, in these terms,

$$I(f) = 2\pi^2 \sum_{n \in \mathbb{Z}} n^2 |a_n|^2. \quad (6)$$

The formula (4) for signed area translates to

$$A(f) = \pi \sum_{n \in \mathbb{Z}} n |a_n|^2 = \pi \sum_{n=1}^{\infty} n (|a_n|^2 - |a_{-n}|^2), \quad (7)$$

It is clear from (6) and (7) that any $f : [0, 1] \rightarrow \mathbb{C}$ that minimises $I(f)$ among those functions for which $A(f) \geq \pi$, and $f(0) = f(1)$, has $a_n = 0$ for $n < 0$. It also follows from (6) and (7), that

$$I(f) \geq 2\pi A(f), \quad (8)$$

for f having only positive Fourier modes. Thus, we must have $I(f) \geq 2\pi^2$ for functions f such that $A(f) \geq \pi$. Note however that if $f(x) = -a + a \exp\{2\pi i x\}$, for $a \in \mathbb{C}$ such that $|a| = 1$, then $I(f) = 2\pi^2$ and $A(f) = \pi$, so that equality in (8) is attained for such functions f . Noting that, if f has only positive Fourier modes and has some $a_n \neq 0$ for $n > 1$, the inequality in (8) is strict, we have deduced that each of the minimising functions takes the form of a progression at constant rate along the circumference of a circle of radius 1.

However, it appears that the fluctuation behaviour of the conditioned process may not be understood by a direct application of the techniques of large deviations. We now begin to develop the tools required for our study of this deviation.

2.2 Tools for the proof

Firstly, we find a lower bound on the probability that the Brownian loop captures the required area.

Lemma 1 *Let $G_m = G_m(T)$ denote a regular polygon with m vertices that contains an area equal to πT^2 . Then, for any constant C_1 satisfying $C_1 > 4/3$, and for all $T \geq T_{C_1}$ sufficiently high,*

$$\begin{aligned} & \mathbb{P}\left(\text{enc}(B) \supset x + G_{\lfloor T^{\frac{1}{3}} \rfloor} \text{ for some } x \in \mathbb{R}^2\right) \\ & \geq \exp\left\{-2\pi^2 T - C_1 T^{\frac{1}{3}} \log T\right\}. \end{aligned}$$




Figure 3: A realization of the event A_i .

Proof Denote the successive vertices of $G_m(T)$ by

$$\{y_i : i \in \{1, \dots, m\}\},$$

where we set $m = \lfloor T^{1/3} \rfloor$. and let D_i denote the line segment whose end-points are the centre of G_m , and y_i . Let ϕ_i denote the disk of radius one whose centre lies on the continuation of D_i at distance two from y_i . Let M_i denote the open half-plane, disjoint from $G_m(T)$, whose boundary contains the line segment $[y_i, y_{i+1}]$ (in the case where $i = m$, the line segment $[y_m, y_1]$). Let $q \in \mathbb{R}^2$ be such that $q \in -\phi_1$. For $i \in \{0, \dots, m-2\}$, let A_i denote the event that

$$\begin{aligned} B\left(\frac{iT}{m}\right) &\in q + \phi_{i+1}, \\ B\left(\frac{(i+1)T}{m}\right) &\in q + \phi_{i+2}, \end{aligned}$$

and

$$B(t) \in q + M_{i+1} \text{ for } t \in \left[\frac{iT}{m}, \frac{(i+1)T}{m}\right].$$

In the case where $i = m-1$, we use the same definition, with ϕ_1 replacing ϕ_{i+2} in its statement. Note that

$$\bigcap_{i=0}^{m-1} A_i \subseteq \left\{ \text{enc}(B) \supset q + G_m \right\}. \quad (9)$$

We claim that

$$\mathbb{P}\left(\bigcap_{i=1}^m \left\{B\left(\frac{iT}{m}\right) \in q + \phi_{i+1}\right\}\right) \geq \exp\left\{-2\pi^2 T - (2/3 + o(1))T^{\frac{1}{3}} \log T\right\}. \quad (10)$$

To see this, note that the left-hand-side of (10) is given by

$$\frac{m^m}{(2\pi T)^{m-1}} \int \exp\left\{-\frac{m}{2T} \sum_{i=0}^{m-1} \|x_{i+1} - x_i\|^2\right\} dx_1 \dots dx_{m-1}, \quad (11)$$

where the range of integration is equal to $(q + \phi_2) \times \dots \times (q + \phi_m)$ and where we set $x_0 = x_m = 0$. The form for the expression in (11) occurs by computing the density of a finite-dimensional distribution of a Brownian bridge as the ratio of the corresponding density for a Brownian motion and the density at zero of a normal random variable with mean zero and variance T . It is straightforward to show that

$$|D_i| = T + \frac{\pi^2}{3}T^{\frac{1}{3}} + O(1). \quad (12)$$

Thus, the distance between successive vertices satisfies

$$\begin{aligned} d(y_i, y_{i+1}) &= 2|D_i| \sin \frac{\pi}{m} \leq \frac{2|D_i|\pi}{m} \\ &\leq \frac{2\pi T}{m} + O(1). \end{aligned} \quad (13)$$

From (13), it follows that the expression in (11) is bounded below by

$$\frac{m^m}{(2\pi T)^{m-1}} \left[\prod_{i=2}^m \text{vol}(\phi_i) \right] \exp\left\{\frac{-m^2}{2T} \left(\frac{2\pi T}{m} + O(1)\right)^2\right\}.$$

Since $\text{vol}(\phi_i) = \pi$ for $i \in \{1, \dots, \lfloor T^{1/3} \rfloor\}$, this expression is bounded below by $\exp\left\{-2\pi^2 T - (2/3 + o(1))T^{\frac{1}{3}} \log T\right\}$, as required to demonstrate that (10) holds. We claim that

$$\begin{aligned} &\mathbb{P}\left(\bigcap_{i=0}^{m-1} A_i \mid \bigcap_{i=1}^m B\left(\frac{iT}{m}\right) \in q + \phi_{i+1}\right) \\ &\geq \exp\left\{-\left(\frac{2}{3} + o(1)\right)T^{1/3} \log T\right\}. \end{aligned} \quad (14)$$

To see this, note that

$$\begin{aligned} &\mathbb{P}\left(\bigcap_{i=0}^{m-1} A_i \mid \bigcap_{i=1}^m B\left(\frac{iT}{m}\right) \in q + \phi_{i+1}\right) \\ &= \prod_{i=0}^{m-1} \mathbb{P}\left(A_i \mid \left\{B\left(\frac{iT}{m}\right) \in q + \phi_{i+1}\right\} \cap \left\{B\left(\frac{(i+1)T}{m}\right) \in q + \phi_{i+2}\right\}\right). \end{aligned} \quad (15)$$

We condition on, for example,

$$B\left(\frac{T}{m}\right) \in q + \phi_2, B\left(\frac{2T}{m}\right) \in q + \phi_3. \quad (16)$$

Consider the one-dimensional process $\hat{B} : I \rightarrow \mathbb{R}$, $I := [T/m, 2T/m]$, that is the component of B in the direction orthogonal to the line segment $[B(T/m), B(2T/m)]$. Under conditioning on (16), $\hat{B}$ is distributed as a Brownian bridge run to and from two given points, each of which we may insist lies in the interval $[-2, 0]$. Note that the motion of B will remain in the half-plane M_1 during the interval of time I provided that $\hat{B} \leq 1/2$ throughout this time. The probability that the maximum of a one-dimensional Brownian bridge that is run for a given time, starts at $a \in \mathbb{R}$ and ends at $b \in \mathbb{R}$, does not exceed a given value is a decreasing function of a and of b . We may assume therefore that $\hat{B}(T/m) = 0$ and $\hat{B}(2T/m) = 0$.

Recall that, if M^+ denotes the maximum of the one-dimensional Brownian bridge run for time T , then, for any $r > 0$,

$$\mathbb{P}(M^+ > r) = \exp\left\{-\frac{2r^2}{T}\right\}. \quad (17)$$

This assertion appears as formula (3.40) in [5, Chapter 4].

We deduce that

$$\mathbb{P}\left(\hat{B}(t) \leq 1/2 \text{ for all } t \in I\right) \geq 1 - \exp\left\{-\frac{m}{2T}\right\} \geq \frac{1}{8}T^{-\frac{2}{3}},$$

the latter inequality being valid for high values of T .

Thus, given the occurrence of (16), the probability that the event A_1 occurs is at least $\frac{1}{8}T^{-2/3}$. From this, and the product form of (15), follows (14).

From (10) and (14), we find that

$$\mathbb{P}\left(\bigcap_{i=0}^{m-1} A_i\right) \geq \exp\left\{-2\pi^2 T - \left(\frac{4}{3} + o(1)\right)T^{\frac{1}{3}} \log T\right\}.$$

The statement of the lemma follows from the inclusion (9). $\square$

The following result is required.

Lemma 2 *For any constant $C_2 \geq 128\pi$, the planar Brownian loop $B : [0, T] \rightarrow \mathbb{R}^2$ has the property that*

$$\sup_{t \in [0, T]} |B(t)|^2$$

is bounded above in distribution by $C_2 T + Z^2$, where Z is a normal random variable with mean zero and variance T . That is, for all $a \in (0, \infty)$,

$$\mathbb{P}\left(\sup_{t \in [0, T]} |B(t)|^2 \geq a\right) \leq \mathbb{P}\left(C_2 T + Z^2 \geq a\right).$$

Remark In fact, the method of proof we give would permit us to pick Z to be distributed normally with mean zero and variance $(1/2 + \epsilon)T$, for any $\epsilon > 0$ (provided that the constant C_2 is changed suitably.) We made the choice $\epsilon = 1/2$, because it is not valuable in the application to choose any lower value for ϵ .

Proof Note that

$$\sup_{t \in [0, T]} |B(t)|^2 \leq \sup_{t \in [0, T]} B_1(t)^2 + \sup_{t \in [0, T]} B_2(t)^2, \quad (18)$$

where B_1 and B_2 denote the components of B . From the probability (17) that the maximum of a one-dimensional Brownian bridge exceeds a given level, and its counterpart for the minimum value reaching below a prescribed value, it follows that

$$\mathbb{P}\left(\sup_{t \in [0, T]} B_j(t)^2 > r^2\right) \leq 2 \exp\left\{-\frac{2r^2}{T}\right\}, \text{ for } j \in \{1, 2\}. \quad (19)$$

Replacing r by $r/\sqrt{2}$ in (19), we obtain from (18) that

$$\mathbb{P}\left(\sup_{t \in [0, T]} |B(t)|^2 > r^2\right) \leq 4 \exp\left\{-\frac{r^2}{T}\right\}. \quad (20)$$

A lower bound on the tail of a random variable Z , distributed normally with mean zero and variance T , is now obtained:

$$\begin{aligned} \mathbb{P}(Z^2 > r^2) &= \mathbb{P}\left(\frac{|Z|}{\sqrt{T}} > \frac{r}{\sqrt{T}}\right) \\ &\geq \frac{\frac{2r}{\sqrt{T}}}{\left(\frac{r^2}{T} + 1\right)\sqrt{2\pi}} \exp\left\{-\frac{r^2}{2T}\right\}, \end{aligned} \quad (21)$$

where the inequality follows from a standard bound on the tail of the normal distribution, presented in Section 14.8 of [9]. From (20) and (21), it follows that

$$\mathbb{P}\left(\sup_{t \in [0, T]} |B(t)|^2 > r^2\right) \leq \mathbb{P}(Z^2 > r^2),$$

for $r \geq C\sqrt{T}$, where $C = 8\sqrt{2\pi}$. We have derived

$$\mathbb{P}\left(\sup_{t \in [0, T]} |B(t)|^2 \geq a\right) \leq \mathbb{P}\left(C_2 T + Z^2 \geq a\right). \quad (22)$$

for any $C_2 \geq 0$ and each $a \geq C^2 T$. By choosing $C_2 = C^2$, the right-hand-side (22) becomes equal to 1 for any $a \in [0, C^2 T]$. This establishes the statement of the Lemma. $\square$

We require some notation before proceeding.

Definition 1 Let $m \in \mathbb{N}$ and let $t' \in [0, T]$.

- Let $P = P_m^{t'}$ denote the polygon whose vertices are given by

$$\left\{ B\left(\frac{jT}{m} + t'\right) : j \in \{0, \dots, m-1\} \right\}.$$

- Let the length of the edges of $P_m^{t'}$ be denoted by $\{L_i : i \in \{1, \dots, m\}\}$, so that

$$L_i = \left| B\left(\frac{iT}{m} + t'\right) - B\left(\frac{(i-1)T}{m} + t'\right) \right|,$$

and let l_i denote the line segment whose endpoints are $B((i-1)T/m + t')$ and $B(iT/m + t')$.

- for $i \in \{1, \dots, m\}$, let $R_i = R_i(t')$ denote the maximum of the distance of the range of the motion B from the segment l_i during that interval of time in which l_i is traversed by B . This is,

$$R_i = \sup_{t \in [0, T/m]} d\left(B\left(\frac{(i-1)T}{m} + t' + t\right), l_i\right),$$

where d denotes the distance between two sets in $\mathbb{R}^2$.

- for $i \in \{1, \dots, m\}$, let $\hat{R}_i = \hat{R}_i(t')$ denote the maximum of the absolute value of the displacement of the motion from a point that traverses l_i at a linear rate during the time $[(i-1)T/m, iT/m]$. That is,

$$\begin{aligned} \hat{R}_i &= \sup_{t \in [0, T/m]} d\left(B\left(\frac{(i-1)T}{m} + t' + t\right), \right. \\ &\quad \left. \left(1 - \frac{mt}{T}\right)B\left(\frac{(i-1)T}{m} + t'\right) + \frac{mt}{T}B\left(\frac{iT}{m} + t'\right)\right). \end{aligned}$$

- let the set $Q_i = Q_i(t')$ be given by $Q_i = \{x \in \mathbb{R}^2 : d(x, l_i) \leq R_i\}$.

Lemma 3 For any $T \in (0, \infty)$, $t' \in [0, T/m]$ and $m \in \mathbb{N}$,

$$\text{enc}(B) \subseteq \text{conv}(P) \cup \bigcup_{i=1}^m Q_i. \quad (23)$$

Proof The objects used in the proof are depicted in Figure 3. Take a point $y \in \text{enc}(B) \setminus \text{conv}(P)$. Locate a half-plane H that contains $\text{conv}(P)$ and excludes y . Let l' denote the line through y that intersects the boundary of H at right angles. Note that there exists a point z in the range of B that lies on l' , at a distance from the half-plane H greater than that of y : this is because $y \in \text{enc}(B)$. Let l_i be the interval in the polygon P traversed while




Figure 4: The objects of Definition 1




Figure 5: The proof of Lemma 3

the Brownian path captures the point z : that is $i \in \{1, \dots, m\}$ is chosen so that

$$z = B(t), \text{ where } t \in \left[\frac{(i-1)T}{m} + t', \frac{iT}{m} + t' \right]. \quad (24)$$

The point y is closer to any given point in H than is z , and each point in the line segment l_i lies in $\text{conv}(P)$, and so in H . Hence, $d(y, l_i) \leq d(z, l_i)$. That $d(z, l_i) \leq R_i$ follows from (24). Hence, $y \in Q_i$. $\square$

Lemma 4 *Let $\epsilon \in [0, 1/6]$, and let $m = \lfloor T^{1/3-\epsilon} \rfloor$. Suppose that the point $t' \in [0, T]$ in the definition of the polygon P is fixed, or that it is sampled randomly, according to an arbitrary distribution on $[0, T]$, and independently of the randomness that generates $B : [0, T] \rightarrow \mathbb{R}^2$. For $\epsilon' \in (0, 1/3)$, let $H = H(\epsilon')$ denote the event that*

$$|\text{conv}(P)| \geq \pi T^2 - T^{4/3+\epsilon'},$$

where $\text{conv}(P)$ denotes the convex hull of the polygon P . Provided that $\epsilon' > 2\epsilon$, for sufficiently large values of T ,

$$\mathbb{P}\left(H(\epsilon')^c \cap \{|\text{enc}(B)| \geq \pi T^2\}\right) \leq \exp\left\{-2\pi^2 T - c_1 T^{\frac{1}{3}+2(\epsilon'-\epsilon)}\right\},$$

where c_1 is any positive constant at most $\frac{2^{-10}}{108 \cdot (128)^2 \pi^4}$.

Proof For $a \in \mathbb{N}$, let S_a denote the event that

$$\pi T^2 - 2^a T^{4/3+\epsilon'} > |\text{conv}(P)| \geq \pi T^2 - 2^{a+1} T^{4/3+\epsilon'}.$$

Note that

$$H^c = \left(\bigcup_{a=0}^k S_a \right) \cup R, \quad (25)$$

where $k = \lfloor \log_2 T^{2/3-\epsilon'} \rfloor - 1$, with R denoting the event that the area of $\text{conv}(P)$ is at most $(\pi - 1/2)T^2$.

Note that

$$\begin{aligned} \left| \text{enc}(B) \setminus \text{conv}(P) \right| &\leq \sum_{i=1}^m |Q_i| \\ &\leq 2 \sum_{i=1}^m (L_i + 2R_i) R_i \\ &\leq 2 \sum_{i=1}^m (L_i + 2\hat{R}_i) \hat{R}_i \\ &\leq 2 \left(\sum_{i=1}^m L_i^2 \right)^{1/2} \left(\sum_{i=1}^m \hat{R}_i^2 \right)^{1/2} + 4 \sum_{i=1}^m \hat{R}_i^2, \end{aligned} \quad (26)$$

where the first inequality follows from Lemma 3. The second inequality follows from the fact that Q_i is contained in a rectangle of length $L_i + 2R_i$ and width $2R_i$, while the third is implied by the inequality $R_i \leq \hat{R}_i$.

Let $a \in \{0, \dots, k\}$. Note that, on the event $S_a \cap \{|\text{enc}(B)| \geq \pi T^2\}$, we have that

$$\left| \text{enc}(B) \setminus \text{conv}(P) \right| \geq |\text{enc}(B)| - |\text{conv}(P)| \geq 2^a T^{4/3+\epsilon'}. \quad (27)$$

We may write

$$L_i^2 = \frac{T(E_{2i-1}^2 + E_{2i}^2)}{m}, \quad (28)$$

where, for each $i \in \{1, \dots, m\}$, the quantities $\sqrt{T/m}E_{2i-1}$ and $\sqrt{T/m}E_{2i}$ are the horizontal and vertical components of the vector $B(iT/m + t') - B((i-1)T/m + t')$. As such, the family $\{E_i : i \in \{1, \dots, 2m\}\}$ is a collection of independent standard normal random variables, conditioned by insisting that

$$\sum_{i=1}^m E_{2i-1} = 0, \quad \sum_{i=1}^m E_{2i} = 0. \quad (29)$$

For each $i \in \{1, \dots, m\}$ and any pair $(x, y) \in \mathbb{R}^2$, the conditional distribution of the random variable $\hat{R}_i$ given the event that $E_{2i-1} = x$ and $E_{2i} = y$ is independent of x and y . Indeed, the process $Z_i : [(i-1)T/m, iT/m] \rightarrow \mathbb{R}^2$ given by

$$Z_i(t) = B(t) + \left(1 - (mt/T - (i-1))\right) B\left(\frac{(i-1)T}{m}\right) + (mt/T - (i-1)) B\left(\frac{iT}{m}\right)$$

is a standard Brownian loop run for a time of T/m , no matter how we condition the values of the endpoints $B((i-1)T/m)$ and $B(iT/m)$. This means that, under any conditioning of the form $E_{2i-1} = x$ and $E_{2i} = y$, each $\hat{R}_i$ has the distribution of the maximal Euclidean distance of this Brownian loop. As such, we may apply Lemma 2 to find a collection of standard normal random variables $\{F_i : i \in \{1, \dots, m\}\}$ for which

$$\hat{R}_i^2 \leq \frac{T}{m}(C_2 + F_i^2). \quad (30)$$

The fact that the conditional distribution of $\hat{R}_i$ does not depend on x and y implies that we may assume that each F_i is independent of F_j , for $j \in \{1, \dots, m\}$ with $j \neq i$, and of each E_j , for $j \in \{1, \dots, 2m\}$.

From (26), (27), (28) and (30) follows

$$\frac{2T}{m} \left(\sum_{i=1}^{2m} E_i^2 \right)^{1/2} \left(C_2 m + \sum_{i=1}^m F_i^2 \right)^{1/2} + 4C_2 T + \frac{4T}{m} \sum_{i=1}^m F_i^2 \geq 2^a T^{4/3+\epsilon'}.$$

Using the inequalities $m \geq T^{1/3-\epsilon}/2$ and $C_2 \geq \sqrt{C_2} \geq 1$,

$$\left(\sum_{i=1}^{2m} E_i^2\right)^{1/2} \left(m + \sum_{i=1}^m F_i^2\right)^{1/2} + 2m + 2 \sum_{i=1}^m F_i^2 \geq \frac{2^{a-2} T^{2/3-\epsilon+\epsilon'}}{C_2}. \quad (31)$$

We note that

$$\sum_{i=1}^m L_i^2 \geq \frac{(\sum_{i=1}^m L_i)^2}{m} \geq \frac{(\text{arcl}(\text{conv}(P)))^2}{m} \geq \frac{4\pi|\text{conv}(P)|}{m}, \quad (32)$$

where arcl denotes arclength. The successive inequalities in (32) are consequences of the Cauchy-Schwarz inequality, the fact that the arclength of any polygon exceeds that of its convex hull, and the standard isoperimetric inequality, which in this case asserts that

$$(\text{arcl}(\text{conv} P))^2 \geq 4\pi|\text{conv}(P)|.$$

It follows from (28),(32) and the lower bound on $|\text{conv}(P)|$ provided by the occurrence of the event S_a that

$$\sum_{i=1}^{2m} E_i^2 \geq 4\pi^2 T - 2^{a+3} \pi T^{1/3+\epsilon'}, \quad (33)$$

because the left and right-hand-side of this inequality are respectively bounded below and above by the quantity $\frac{4\pi|\text{conv}(P)|}{T}$. We see that the event

$$\left\{|\text{enc}(B)| \geq \pi T^2\right\} \cap S_a$$

is contained in the event F , specified by the occurrence of the inequalities in (31) and (33). Note that, we have the following inclusion:

$$F \subseteq A_C^1 \cup A_C^2 \cup A_C^3, \quad (34)$$

where the events on the right-hand-side are given by

$$\begin{aligned} A_C^1 &= \left\{ \sum_{i=1}^{2m} E_i^2 \geq CT \right\}, \\ A_C^2 &= \left\{ \sum_{i=1}^{2m} E_i^2 \in [4\pi^2 T - 2^{a+3} \pi T^{1/3+\epsilon'}, CT) \right\} \\ &\cap \left\{ m + \sum_{i=1}^m F_i^2 \geq \sum_{i=1}^{2m} E_i^2 \right\} \end{aligned}$$

and

$$A_C^3 = \left\{ \sum_{i=1}^{2m} E_i^2 \in [4\pi^2 T - 2^{a+3}\pi T^{1/3+\epsilon'}, CT) \right\} \\ \cap \left\{ m + \sum_{i=1}^m F_i^2 \geq \frac{2^{2a-4}T^{1/3-2\epsilon+2\epsilon'}}{9CC_2^2} \right\}.$$

Here, C denotes a fixed constant, and the constant C_2 satisfies the bound stated in Lemma 2. To derive (34), note that, if the event $F \cap (A_C^1 \cup A_C^2)^c$ occurs, then

$$3 \left(\sum_{i=1}^{2m} E_i^2 \right)^{\frac{1}{2}} \left(m + \sum_{i=1}^m F_i^2 \right)^{\frac{1}{2}} \\ \geq \left(\sum_{i=1}^{2m} E_i^2 \right)^{\frac{1}{2}} \left(m + \sum_{i=1}^m F_i^2 \right)^{\frac{1}{2}} + 2 \left(m + \sum_{i=1}^m F_i^2 \right) \\ \geq \frac{2^{a-2}}{C_2} T^{\frac{2}{3}-\epsilon+\epsilon'}.$$
(35)

The first inequality follows from the fact that $m + \sum_{i=1}^m F_i^2 < \sum_{i=1}^{2m} E_i^2$, which holds on the event $F \cap (A_C^1 \cup A_C^2)^c$, while the second is simply (31). Using (35), and the fact that $\sum_{i=1}^{2m} E_i^2 < CT$, we find that the second event of the two whose intersection defines A_C^3 must occur. The first occurs provided that both the event $(A_C^1)^c$ occurs and the inequality (33) is satisfied, which means that it occurs if the event $F \cap (A_C^1 \cup A_C^2)^c$ does.

The standard normal random variables $\{E_i : i \in \{1, \dots, 2m\}\}$ are independent, conditioned only by the two linear constraints (29). Thus, the quantity $\sum_{i=1}^{2m} E_i^2$ has the χ^2 -distribution with $2m - 2$ degrees of freedom. The random variable $\sum_{i=1}^m F_i^2$ has the χ^2 -distribution with m degrees of freedom. Recall that the density $f_m : [0, \infty) \rightarrow [0, \infty)$ of this latter distribution is given by

$$f_m(x) = \frac{x^{m/2-1} \exp\{-x/2\}}{2^{m/2}\Gamma(m/2)}.$$
(36)

From this formula, we find that, for any given $C > 0$ and all sufficiently high values of T ,

$$\mathbb{P}(A_C^1) \leq \exp\left\{-\frac{CT}{4}\right\}.$$
(37)

Recalling the independence property of the family of random variables $\{E_i : i \in \{1, \dots, 2m\}\}$ and $\{F_i : i \in \{1, \dots, m\}\}$ asserted after (30), it follows that

$$\mathbb{P}(A_C^2) \leq \int_l^{CT} f_{2m-2}(x) dx \int_{l-m}^{\infty} f_m(x) dx,$$
(38)

where $l = 4\pi^2 T - 2^{a+3} \pi T^{1/3+\epsilon'}$. The right-hand-side of (38) may be bounded above by

$$CT(CT)^{T^{1/3-\epsilon}} \exp \left\{ -2\pi^2 T + 2^{a+2} \pi T^{\frac{1}{3}+\epsilon'} \right\} \times \\ \left[CT(CT)^{T^{1/3-\epsilon}} \exp \left\{ -2\pi^2 T + 2^{a+2} \pi T^{\frac{1}{3}+\epsilon'} + \frac{1}{2} T^{\frac{1}{3}-\epsilon} \right\} + \exp \left\{ -CT/4 \right\} \right]$$

by making use of $m \leq T^{1/3-\epsilon}$. From the fact that $2^{a+1} \leq T^{2/3-\epsilon'}$ follows that for any constant c' and for all T exceeding a c' -dependent constant,

$$\mathbb{P}(A_C^2) \leq \exp \left\{ - \left[4\pi(\pi - 1) - c' \right] T \right\}, \quad (39)$$

provided that $C > 8\pi(\pi - 1)$. On the event A_3 , $\sum_{i=1}^m F_i^2 \geq m$: this follows from the second requirement in the definition of A_3 , along with the weaker than given inequality $\epsilon < 2\epsilon'$, and $m \leq T^{1/3-\epsilon}$. Thus, if A_3 occurs, then

$$\sum_{i=1}^m F_i^2 \geq \frac{2^{2a-5} T^{1/3-2\epsilon+2\epsilon'}}{9CC_2^2}.$$

Thus, by a similar estimate as in the case of A_C^2 ,

$$\mathbb{P}(A_C^3) \leq CT(CT)^{T^{1/3-\epsilon}} \exp \left\{ -2\pi^2 T + 2^{a+2} \pi T^{1/3+\epsilon'} \right\} \\ \times \left[CT(CT)^{T^{1/3-\epsilon}} \exp \left\{ -\frac{2^{2a-6}}{9CC_2^2} T^{\frac{1}{3}-2\epsilon+2\epsilon'} \right\} + \exp \left\{ -CT/4 \right\} \right].$$

Using the fact that $\epsilon' > 2\epsilon$,

$$\mathbb{P}(A_C^3) \leq \exp \left\{ -2\pi^2 T - \frac{2^{2a-7}}{9CC_2^2} T^{\frac{1}{3}-2\epsilon+2\epsilon'} \right\} + \exp \left\{ -CT/5 \right\}. \quad (40)$$

We find that

$$\mathbb{P} \left(S_a \cap \left\{ |\text{enc}(B)| \geq \pi T^2 \right\} \right) \\ \leq \mathbb{P} \left(A_C^1 \cup A_C^2 \cup A_C^3 \right) \leq \mathbb{P}(A_C^1) + \mathbb{P}(A_C^2) + \mathbb{P}(A_C^3) \\ \leq \exp \left\{ -2\pi^2 T - \frac{2^{2a-8}}{9CC_2^2} T^{\frac{1}{3}-2\epsilon+2\epsilon'} \right\} + \exp \left\{ -CT/6 \right\}$$

where in the first inequality, we used (34), and in the third, (37), (39) and (40). Recalling that $k = \lfloor \log_2 T^{2/3-\epsilon'} \rfloor - 1$, we find that, provided that $C > 12\pi^2$,

$$\mathbb{P} \left(\left(\bigcup_{a=0}^k S_a \right) \cap \left\{ |\text{enc}(B)| \geq \pi T^2 \right\} \right) \leq \exp \left\{ -2\pi^2 T - c_1 T^{\frac{1}{3}-2\epsilon+2\epsilon'} \right\},$$

for any constant c_1 satisfying

$$c_1 \in \left(0, \frac{2^{-9}}{9CC_2^2}\right) \subseteq \left(0, \frac{2^{-10}}{9 \cdot 12 \cdot (128)^2 \pi^4}\right),$$

where the second inclusion is ensured by choosing $C_2 = 128\pi$ and C slightly above $12\pi^2$.

By (25), it only remains to bound the probability of the event R that appears in there. Note that if R occurs, then

$$|\text{enc}(B) \setminus \text{conv}(P)| \geq \frac{1}{2}T^2.$$

From (26), we see that the occurrence of R implies that

$$2\left(\sum_{i=1}^m L_i^2\right)^{1/2} \left(\sum_{i=1}^m \hat{R}_i^2\right)^{1/2} + 4\sum_{i=1}^m \hat{R}_i^2 \geq \frac{1}{2}T^2,$$

and so that

$$\left(\sum_{i=1}^{2m} E_i^2\right)^{1/2} \left(C_2m + \sum_{i=1}^m F_i^2\right)^{1/2} + 2\left(C_2m + \sum_{i=1}^m F_i^2\right) \geq \frac{1}{8}T^{\frac{4}{3}-\epsilon}, \quad (41)$$

by use of (28), (30) and the inequality $m \geq \frac{1}{2}T^{1/3-\epsilon}$. If the inequality (41) is satisfied, then so is one of

$$C_2m + \sum_{i=1}^m F_i^2 \geq \frac{1}{24}T^{\frac{4}{3}-\epsilon}$$

and

$$\sum_{i=1}^{2m} E_i^2 \geq \frac{1}{24}T^{\frac{4}{3}-\epsilon}.$$

Since $m \leq T^{1/3}$ and $\epsilon < 1/3$, each of these inequalities is satisfied with a probability that decays at a rate faster than exponential. This completes the proof. $\square$

It is of interest to bound the extent of the excess of area captured by the motion, partly for the reasons presented in the heuristic discussion that ends the Introduction. In addition, we will make use of the following bound in the proof of Theorem 1.

Proposition 1 *For $\delta \in (0, 1/3)$ and for all $T \in (0, \infty)$ sufficiently high, we have that*

$$\mathbb{P}\left(|\text{enc}(B)| \geq \pi T^2 + T^{\frac{4}{3}+\delta}\right) \leq \exp\left\{-2\pi^2 T - \frac{\pi}{4}T^{\frac{1}{3}+\delta}\right\}.$$

Proof We set $m = \lfloor T^{\frac{1}{3}} \rfloor$. Note that the event whose probability we seek to bound lies in $Y_1 \cup Y_2$, where the events Y_1 and Y_2 are given by

$$Y_1 = \left\{ |\text{enc}(B)| \geq \pi T^2 + T^{\frac{4}{3}+\delta} \right\} \cap \left\{ |\text{conv}(P)| < \pi T^2 + \frac{1}{2} T^{\frac{4}{3}+\delta} \right\},$$

and by

$$Y_2 = \left\{ |\text{conv}(P)| \geq \pi T^2 + \frac{1}{2} T^{\frac{4}{3}+\delta} \right\}.$$

Let $\hat{T} \in (0, \infty)$ be given by $\pi \hat{T}^2 = \pi T^2 + T^{\frac{4}{3}+\delta}$. We find that

$$\begin{aligned} \mathbb{P}(Y_1) &\leq \mathbb{P}\left(\left\{ |\text{enc}(B)| \geq \pi \hat{T}^2 \right\} \cap \left\{ |\text{conv}(P)| < \pi \hat{T}^2 - \frac{1}{4} \hat{T}^{\frac{4}{3}+\delta} \right\}\right) \quad (42) \\ &\leq \exp\left\{ -2\pi^2 T - \frac{c_1}{4^2} T^{\frac{1}{3}+2\delta} \right\}. \end{aligned}$$

The first inequality in (42) follows for high values of T from the fact that $\delta < 2/3$. The second is an application of Lemma 4 (with the choice $\epsilon' = \delta - \frac{\log 4}{\log \hat{T}}$), and of the inequality $\hat{T} \geq T$.

Note that, by (28) and (32),

$$\sum_{i=1}^{2m} E_i^2 \geq \frac{4\pi}{T} |\text{conv}(P)|,$$

so that

$$\mathbb{P}(Y_2) \leq \mathbb{P}\left(\sum_{i=1}^{2m} E_i^2 \geq 4\pi^2 T + 2\pi T^{\frac{1}{3}+\delta}\right) \leq \exp\left\{ -2\pi^2 T - \frac{\pi}{2} T^{\frac{1}{3}+\delta} \right\}. \quad (43)$$

We used the form (36) of the density of the χ^2 -distribution in the second inequality, as well as the fact that $m \leq T^{1/3}$. Applying the bounds (42) and (43) to estimate $\mathbb{P}(Y_1 \cup Y_2)$, we deduce the statement of the proposition. $\square$

Lemma 5 *Given functions $f, g : [0, T] \rightarrow [0, \infty)$, let $Q_{f,g}$ denote the event that*

$$\sup_{t \in [0, T]} \sup_{s \in [t, t+f(T)]} |B(s) - B(t)| \leq g(T).$$

Provided that $f(T) \leq T$, we have that

$$\mathbb{P}(Q_{f,g}^c) \leq \left(\frac{32\sqrt{2}T}{\sqrt{\pi f(T)g(T)}} + \frac{16\sqrt{T}}{\sqrt{\pi}g(T)} \right) \exp\left\{ -\frac{g(T)^2}{128f(T)} \right\}. \quad (44)$$

Proof: Firstly, note that

$$Q_{f,g}^c \subseteq \bigcup_{j=1}^{\lfloor T/f(T) \rfloor} C_j, \quad (45)$$

where C_j is the event that there exist $t_1, t_2 \in [jf(T), (j+2)f(T)]$ for which $|B(t_1) - B(t_2)| > g(T)$. Noting that the process $B_t : [0, T] \rightarrow \mathbb{R}^2$ given by $B_t(s) = B(t+s) - B(t)$ has the same law as $B = B_0$, we see that

each of the events C_j for $j \in \{1, \dots, \lfloor T/f(T) \rfloor\}$ has an equal probability. (46)

We represent $B : [0, T] \rightarrow \mathbb{R}^2$ in the form $B(t) = W(t) - tW(T)/T$, where $W : [0, T] \rightarrow \mathbb{R}^2$ is distributed as a standard planar Brownian motion. Note then the inclusion

$$C_1 \subseteq (A_1 \cap F^c) \cup F, \quad (47)$$

where A_1 is the event that there exist $t_1, t_2 \in [f(T), 3f(T)]$ for which $|W(t_1) - W(t_2)| > g(T)/2$, and where F is the event

$$F = \left\{ W(T) > \frac{Tg(T)}{4f(T)} \right\}.$$

Indeed, on the event $C_1 \cap F^c$, we find that

$$\begin{aligned} |W(t_1) - W(t_2)| &\geq |B(t_1) - B(t_2)| - \frac{|t_1 - t_2|}{T} |W(T)| \\ &\geq g(T) - |t_1 - t_2| \frac{g(T)}{4f(T)} \geq \frac{1}{2} g(T), \end{aligned}$$

where $t_1, t_2 \in [0, T]$ are as in the definition of the event C_j . Thus, A_j occurs for that j for which $t_1, t_2 \in [jf(T), (j+2)f(T)]$. We have verified (47).

Given that, for each $j \in \{1, \dots, \lfloor T/f(T) \rfloor\}$,

$$\mathbb{P}(C_j) = \mathbb{P}(C_1) \leq \mathbb{P}(A_1) + \mathbb{P}(F), \quad (48)$$

we seek to bound the probability of the events A_1 and F . To this end, note that, if A_1 occurs, then one of the components in the x or y direction of $W(t_1) - W(t_2)$ on $[f(T), 3f(T)]$ has a difference between its maximum and minimum values that exceeds $g(T)/(2\sqrt{2})$. For this component, one of the maximum and the absolute value of the minimum exceeds $g(T)/(4\sqrt{2})$. Thus,

$$\begin{aligned} \mathbb{P}(A_1) &\leq 4\mathbb{P}\left(\sup_{t \in [0, 2f(T)]} W_1(t) > \frac{g(T)}{4\sqrt{2}}\right) \\ &= 8\mathbb{P}\left(W_1(2f(T)) > \frac{g(T)}{4\sqrt{2}}\right) \\ &\leq \frac{32\sqrt{2}\sqrt{f(T)}}{\sqrt{\pi}g(T)} \exp\left\{-\frac{g(T)^2}{128f(T)}\right\}, \end{aligned} \quad (49)$$

where in the equality, we used the reflection principle, and, in the latter inequality, a standard tail bound for a normal random variable (Theorem 1.4 of [4]).

To bound the probability of the event F , note that if $\frac{f(T)}{T}|W(T)| > \frac{1}{4}g(T)$, then at least one of the inequalities

$$\frac{f(T)}{T}|W_i(T)| > \frac{1}{4\sqrt{2}}g(T).$$

holds for $i \in \{1, 2\}$. By the same tail bound,

$$\begin{aligned} \mathbb{P}(F) &= \mathbb{P}\left(\frac{f(T)}{T}|W(T)| > \frac{1}{4}g(T)\right) \\ &\leq \frac{16f(T)}{\sqrt{\pi T}g(T)} \exp\left\{-\frac{Tg(T)^2}{64f(T)^2}\right\} \leq \frac{16f(T)}{\sqrt{\pi T}g(T)} \exp\left\{-\frac{g(T)^2}{64f(T)}\right\}, \end{aligned} \quad (50)$$

since $f(T) \leq T$.

We now find, as required, that

$$\begin{aligned} \mathbb{P}(Q_{f,g}^c) &\leq \frac{T}{f(T)}\mathbb{P}(C_1) \leq \frac{T}{f(T)}(\mathbb{P}(A_1) + \mathbb{P}(F)) \\ &\leq \left(\frac{32\sqrt{2}T}{\sqrt{\pi}\sqrt{f(T)}g(T)} + \frac{16\sqrt{T}}{\sqrt{\pi}g(T)}\right) \exp\left\{-\frac{g(T)^2}{128f(T)}\right\}, \end{aligned}$$

by means of (45), (46) in the first inequality, (48) in the second, and (49) and (50) in the third. $\square$

Proposition 2 For $\epsilon_0 \in (0, 1/3)$, let $\overline{J}_{\epsilon_0}$ denote the event that, for some $\epsilon \in (\epsilon_0, 1/3)$ and for some $t \in [0, T]$,

$$\left|B(t + T^{2/3+\epsilon}) - B(t)\right| \geq T^{\frac{2}{3}+2\epsilon}.$$

Then

$$\mathbb{P}\left(\overline{J}_{\epsilon_0} \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\}\right) \leq C_3 T^2 \log T \exp\left\{-2\pi^2 T - 2^{-9} T^{\frac{2}{3}+3\epsilon_0}\right\},$$

where C_3 is any constant exceeding $256\pi^2$.

Lemma 6 For each $\epsilon_0 \in (0, 1/3)$ and $\delta \in (0, \epsilon_0)$, the occurrence of $\overline{J}_{\epsilon_0-\delta}^c$ implies that, for each $\epsilon > \epsilon_0$, if $t, t' \in [0, T]$ satisfy $|B(t') - B(t)| \geq \frac{9}{2}T^{2/3+2\epsilon}$, then $|t' - t| \geq \frac{1}{2}T^{2/3+\epsilon}$.

Proof Given ϵ, t and t' satisfying the hypotheses of the lemma, we find that either

$$\left|B\left(t + T^{\frac{2}{3}+\epsilon}\right) - B(t)\right| \geq \frac{9}{4}T^{\frac{2}{3}+2\epsilon}, \quad (51)$$

or

$$\left|B\left(t + T^{\frac{2}{3}+\epsilon}\right) - B(t')\right| \geq \frac{9}{4}T^{\frac{2}{3}+2\epsilon}. \quad (52)$$

If (51) applies, then certainly $\overline{J}_{\epsilon_0}$ occurs, since $\epsilon > \epsilon_0$. This means of course that $\overline{J}_{\epsilon_0 - \delta}$ occurs as well. Supposing the second eventuality (52) and that $|t' - t| < \frac{1}{2}T^{2/3+\epsilon}$, we note that (52) may be rewritten

$$\left| B\left(t' + T^{\frac{2}{3}+\epsilon'}\right) - B(t') \right| \geq \frac{9}{4}T^{\frac{2}{3}+2\epsilon}, \quad (53)$$

where the quantity ϵ' is easily shown to satisfy the bounds

$$\epsilon' \in (\epsilon - \log 2 / \log T, \epsilon + \log(3/2) / \log T) \quad (54)$$

and thus $\epsilon' > \epsilon - \delta > \epsilon_0 - \delta$, for high values of T . The right-hand-side of (53) exceeds $T^{2/3+2\epsilon'}$ for high values of T , because (54) implies that $T^{2(\epsilon-\epsilon')} \geq 4/9$. We have shown that the occurrence of $\overline{J}_{\epsilon_0 - \delta}$ is a consequence of (52) and the inequality $|t' - t| < \frac{1}{2}T^{2/3+\epsilon}$. This establishes the statement of the lemma. $\square$

The proof of Proposition 2 depends largely on the following Lemma.

Lemma 7 *Let $\epsilon \in (0, 1/3)$. Let $J = J(\epsilon)$ denote the event that, for some $t \in [0, T]$,*

$$\left| B(t + T^{2/3+\epsilon}) - B(t) \right| \geq \frac{1}{2}T^{\frac{2}{3}+2\epsilon}.$$

Then

$$\mathbb{P}\left(J \cap \left\{ |\text{enc}(B)| \geq \pi T^2 \right\}\right) \leq \exp \left\{ -2\pi^2 T - 2^{-9} T^{\frac{2}{3}+3\epsilon} \right\}.$$

Proof Suppose that $\epsilon < 1/5$ (the other case is simple, and will be handled at the end of the proof). Find $\epsilon' > 0$ satisfying $\epsilon' < 2/3$, $\epsilon' - 3\epsilon < 1/3$ and $5\epsilon + 1/3 < 2\epsilon'$ (these conditions may be satisfied, since $\epsilon < 1/5$). We will consider the polygonal approximation of the range of B in the case where $m = \lfloor T^{1/3-\epsilon} \rfloor$. In this context, the quantity t' of Definition 1 will be chosen to be a uniform random variable on the interval $[0, T/m]$ that is independent of the motion B . Note that

$$H(\epsilon') \subseteq \left\{ \sum_{i=1}^m L_i \geq 2\pi T - \frac{\pi}{3} T^{\frac{1}{3}+\epsilon'} \right\}. \quad (55)$$

Indeed,

$$\begin{aligned} \sum_{i=1}^m L_i &\geq 2\sqrt{\pi} \sqrt{|\text{conv}(P)|} \\ &\geq 2\pi \sqrt{T^2 - \frac{T^{4/3+\epsilon'}}{\pi}} \\ &\geq 2\pi T - \frac{\pi}{3} T^{\frac{1}{3}+\epsilon'}, \end{aligned}$$

where the first inequality uses the bound displayed in (32). The second inequality is valid provided that $H(\epsilon')$ occurs and the third is true for all T sufficiently high, since $\epsilon' < 2/3$ and $3 < \pi$.

For $a \geq 0$, let K_a denote the event that

$$2^{a-1}T^{\frac{2}{3}+2\epsilon} \geq L_1 \geq 2^{a-2}T^{\frac{2}{3}+2\epsilon}.$$

We claim that, for any $a \in \mathbb{N}$,

$$H \cap K_a \subseteq \left\{ \sum_{i=1}^{2m} E_i^2 \geq 4\pi^2 T + 2^{2a-5}T^{\frac{2}{3}+3\epsilon} \right\}, \quad (56)$$

where the collection of random variables $\{E_i : i \in \{1, \dots, 2m\}\}$ was introduced in (28). To derive (56), note that

$$\sum_{i=1}^m L_i^2 \geq L_1^2 + \frac{\left(\sum_{i=2}^m L_i\right)^2}{m-1}. \quad (57)$$

From (57) and (28) follows

$$\sum_{i=1}^{2m} E_i^2 \geq \frac{m}{T} L_1^2 + \frac{\left(\sum_{i=2}^{2m} L_i\right)^2}{T}. \quad (58)$$

Supposing the occurrence of the event $H(\epsilon') \cap K_a$, we find that, for high enough values of T ,

$$\begin{aligned} \sum_{i=1}^{2m} E_i^2 &\geq \frac{T^{1/3-\epsilon} - 1}{T} 2^{2a-4}T^{\frac{4}{3}+4\epsilon} + \frac{1}{T} \left(2\pi T - \frac{\pi}{3}T^{\frac{1}{3}+\epsilon'} - 2^{a-1}T^{\frac{2}{3}+2\epsilon} \right)^2 \\ &\geq 2^{2a-4}T^{\frac{2}{3}+3\epsilon} - 2^{2a-4}T^{\frac{1}{3}+4\epsilon} + 4\pi^2 T - \frac{4\pi^2}{3}T^{\frac{1}{3}+\epsilon'} - 2^{a+1}\pi T^{\frac{2}{3}+2\epsilon} \\ &\geq 4\pi^2 T + 2^{2a-5}T^{\frac{2}{3}+3\epsilon}. \end{aligned}$$

In the first inequality, (55) and (58) were used, while, in the third, the relations $\epsilon < 1/3$ and $\epsilon' - 3\epsilon < 1/3$ were required.

From (56) and the definition of the random variables E_i (with $i \in \{1, \dots, 2m\}$), it follows that

$$\mathbb{P}(K_a \cap H) \leq \int_l^\infty f_{2m-2}(y) dy, \quad (59)$$

where f denotes the density of the χ^2 -distribution specified in (36), and the lower limit of integration l is equal to $4\pi^2 T + 2^{2a-5}T^{\frac{2}{3}+3\epsilon}$. Using the fact that $m \leq T^{\frac{1}{3}}$ in estimating the right-hand-side of (59), we find that

$$\mathbb{P}(K_a \cap H(\epsilon')) \leq \exp \left\{ -2\pi^2 T - 2^{2a-7}T^{\frac{2}{3}+3\epsilon} \right\}. \quad (60)$$

for all choices of T exceeding a value that has no dependence on $a \geq 0$. An extra factor of one-half multiplies the term of $T^{\frac{2}{3}+3\epsilon}$ in the last expression, in order to compensate for the non-leading terms in the expression for the χ^2 -density. By summing over $a \in \mathbb{N}$ in (60), we deduce that, for high values of T ,

$$\mathbb{P}\left(\left\{L_1 \geq \frac{1}{4}T^{\frac{2}{3}+2\epsilon}\right\} \cap H(\epsilon')\right) \leq \exp\left\{-2\pi^2 T - 2^{-8}T^{\frac{2}{3}+3\epsilon}\right\}. \quad (61)$$

Allowing I to denote the collection of those times $t \in [0, T]$ for which

$$\left|B\left(t + T^{\frac{2}{3}+\epsilon}\right) - B(t)\right| \geq \frac{1}{2}T^{\frac{2}{3}+2\epsilon},$$

note that the set I is distributionally invariant under shifts of $[0, T]$, and is non-empty if and only if the event $J(\epsilon)$ occurs. We denote by I_1 the event that there exists $t \in I$ satisfying $|t - t'| \leq 1$, where the difference $t - t'$ is being computed modulo T , as each other such will be. We find that

$$\mathbb{P}\left(I_1 \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\} \middle| J(\epsilon) \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\}\right) \geq \frac{2}{T}. \quad (62)$$

If the event I_1 occurs, we have that

$$\begin{aligned} L_1 &= \left|B\left(\frac{T}{m} + t'\right) - B(t')\right| \\ &\geq \left|B\left(t + T^{\frac{2}{3}+\epsilon}\right) - B(t)\right| - \left|B\left(\frac{T}{m} + t'\right) - B\left(t + T^{\frac{2}{3}+\epsilon}\right)\right| - \left|B(t') - B(t)\right| \end{aligned} \quad (63)$$

where $t \in I$ satisfies $|t - t'| \leq 1$. Writing $f(T) = 2T^{\frac{1}{3}+2\epsilon} + 1$ and $g(T) = \frac{1}{8}T^{\frac{2}{3}+2\epsilon}$, note that for high values of T , on the event $Q_{f,g}$,

$$\left|B\left(\frac{T}{m} + t'\right) - B\left(t + T^{\frac{2}{3}}\right)\right| \leq g(T). \quad (64)$$

This is because the difference between the two arguments of B in (64) satisfies

$$\begin{aligned} \left|\left(\frac{T}{m} + t'\right) - \left(t + T^{\frac{2}{3}}\right)\right| &\leq \left|\frac{T}{\lfloor T^{\frac{1}{3}-\epsilon} \rfloor} - T^{\frac{2}{3}+\epsilon}\right| + |t' - t| \\ &\leq \left|\frac{T}{T^{\frac{1}{3}-\epsilon} - 1} - T^{\frac{2}{3}+\epsilon}\right| + 1 \\ &\leq T^{\frac{2}{3}+\epsilon} \sum_{j=1}^{\infty} T^{-j(\frac{1}{3}-\epsilon)} + 1 \leq f(T), \end{aligned} \quad (65)$$

where the fact that $\epsilon \in (0, 1/3)$ was used in the last inequality. Note also that, on the event $Q_{1,g}$,

$$\left|B(t') - B(t)\right| \leq g(T). \quad (66)$$

From (63), (64), (66) and the fact that $t \in I$, it follows that

$$I_1 \cap Q_{f,g} \cap Q_{1,g} \subseteq \left\{ L_1 \geq \frac{1}{4} T^{\frac{2}{3}+2\epsilon} \right\}.$$

Thus,

$$\begin{aligned} & \mathbb{P}\left(\left\{L_1 \geq \frac{1}{4} T^{\frac{2}{3}+2\epsilon}\right\} \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\}\right) \\ & \geq \mathbb{P}\left(I_1 \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\}\right) - \mathbb{P}(Q_{f,g}^c) - \mathbb{P}(Q_{1,g}^c) \\ & \geq \frac{2}{T} \mathbb{P}\left(J(\epsilon) \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\}\right) \\ & \quad - CT \exp\left\{-cT^{1+2\epsilon}\right\} - CT \exp\left\{-\frac{1}{64 \cdot 128} T^{\frac{4}{3}+4\epsilon}\right\}, \end{aligned} \tag{67}$$

with $c \in (0, 128^{-2})$. In the second inequality, (62) was used, as well as the bound provided by Lemma 5. In applying this bound, we used the fact that $f, g \geq 1$ for T sufficiently large. We find that

$$\begin{aligned} & \mathbb{P}\left(J(\epsilon) \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\}\right) \\ & \leq \frac{T}{2} \left[\mathbb{P}\left(\left\{L_1 \geq \frac{1}{4} T^{\frac{2}{3}+2\epsilon}\right\} \cap H(\epsilon')\right) + \mathbb{P}\left(H(\epsilon')^c \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\}\right) \right] \\ & \quad + CT^2 \exp\left\{-cT^{1+2\epsilon}\right\} \\ & \leq \frac{T}{2} \exp\left\{-2\pi^2 T - 2^{-8} T^{\frac{2}{3}+3\epsilon}\right\} \\ & \quad + \frac{T}{2} \exp\left\{-2\pi^2 T - c_1 T^{\frac{1}{3}+2(\epsilon'-\epsilon)}\right\} + CT^2 \exp\left\{-cT^{1+2\epsilon}\right\}, \end{aligned}$$

where (67) was used in the first inequality, the latter requiring (61) and an application of Lemma 4 (so that we use the inequality $\epsilon' > 2\epsilon$, which follows from the assumption that $2\epsilon' > 5\epsilon + 1/3$.) From the fact that $5\epsilon + 1/3 < 2\epsilon'$ follows

$$\mathbb{P}\left(J(\epsilon) \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\}\right) \leq \exp\left\{-2\pi^2 T - 2^{-9} T^{\frac{2}{3}+3\epsilon}\right\},$$

for T sufficiently high, as required.

There remains the case where $\epsilon \in [1/5, 1/3)$. Note that, for any positive value for ϵ , $J(\epsilon) \subseteq Q_{f,g}^c$, with the choices

$$f(T) = T^{\frac{2}{3}+\epsilon}, \quad g(T) = \frac{1}{4} T^{\frac{2}{3}+2\epsilon}$$

being made. We find that, in this case,

$$\begin{aligned} & \mathbb{P}\left(J(\epsilon) \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\}\right) \leq \mathbb{P}(J(\epsilon)) \leq \mathbb{P}(Q_{f,g}^c) \\ & \leq \frac{64}{\sqrt{\pi}} \left(2\sqrt{2} T^{-\frac{5\epsilon}{2}} + T^{-\frac{1}{6}-2\epsilon}\right) \exp\left\{-\frac{1}{2048} T^{\frac{2}{3}+3\epsilon}\right\}, \end{aligned} \tag{68}$$

where Lemma 5 was applied in the third inequality. Given that $2/3 + 3\epsilon > 1$ for a choice of $\epsilon \in [1/5, 1/3)$, we see that (68) establishes the statement of the lemma for such values of ϵ . $\square$

Proof of Proposition 2 We claim that the following inclusion holds, for sufficiently high values of T :

$$\bigcup_{\epsilon'} \left\{ \exists t \in [0, T] : \left| B\left(t + T^{\frac{2}{3} + \epsilon'}\right) - B(t) \right| \geq T^{\frac{2}{3} + 2\epsilon'} \right\} \cap Q_{\frac{1}{CT}, 1} \subseteq J(\epsilon), \quad (69)$$

where the union on the left-hand-side is taken over values of ϵ' satisfying $|\epsilon' - \epsilon| \leq \frac{T^{-5/3 - \epsilon}}{2C \log T}$. To derive (69), firstly set $V_r = \frac{T^{-5/3 - r}}{2C \log T}$ for $r > 0$, and note that, for sufficiently high values of T , and for given $\epsilon > 0$ and any ϵ' satisfying

$$|\epsilon' - \epsilon| \leq V_\epsilon$$

we have that

$$\left| T^{\frac{2}{3} + \epsilon'} - T^{\frac{2}{3} + \epsilon} \right| \leq \frac{1}{CT}. \quad (70)$$

If the event on the left-hand-side of (69) occurs, there exists ϵ' satisfying $|\epsilon' - \epsilon| \leq T^{-5/3 - \epsilon}/(2C \log T)$ and $t \in [0, T]$ for which

$$\left| B\left(t + T^{\frac{2}{3} + \epsilon'}\right) - B(t) \right| \geq T^{\frac{2}{3} + 2\epsilon'}. \quad (71)$$

Note that, provided that the event on the left-hand-side of (69) occurs,

$$\begin{aligned} & \left| B\left(t + T^{\frac{2}{3} + \epsilon}\right) - B(t) \right| \\ & \geq \left| B\left(t + T^{\frac{2}{3} + \epsilon'}\right) - B(t) \right| - \left| B\left(t + T^{\frac{2}{3} + \epsilon'}\right) - B\left(t + T^{\frac{2}{3} + \epsilon}\right) \right| \\ & \geq T^{\frac{2}{3} + 2\epsilon'} - 1 = T^{\frac{2}{3} + 2\epsilon} + (T^{\frac{2}{3} + \epsilon'} - T^{\frac{2}{3} + \epsilon})(T^{\epsilon'} + T^\epsilon) - 1 \geq \frac{1}{2} T^{\frac{2}{3} + 2\epsilon}, \end{aligned}$$

where the final inequality, valid for high values of T , is due to (70) and $\max\{\epsilon, \epsilon'\} < 1$. In the second inequality, we have used (70), (71) and the occurrence of $Q_{1/CT, 1}$. This establishes the inclusion (69).

Note that

$$\overline{J}_{\epsilon_0} \subseteq \bigcup_{i=1}^N \bigcup_{\epsilon' : |\epsilon' - \epsilon_i| \leq V_{\epsilon_i}} \left\{ \exists t \in [0, T] : \left| B\left(t + T^{\frac{2}{3} + \epsilon'}\right) - B(t) \right| \geq T^{\frac{2}{3} + 2\epsilon'} \right\}, \quad (72)$$

where $\{\epsilon_i : i \in \{1, \dots, N\}\}$ denotes a collection of values, each lying in $(\epsilon_0, 1/3)$, for which

$$\left[\epsilon_0, \frac{1}{3} \right] \subseteq \bigcup_{i=1}^N \left(\epsilon_i - V_{\epsilon_i}, \epsilon_i + V_{\epsilon_i} \right).$$

As such, we may choose

$$N \leq CT^2 \log T. \quad (73)$$

From (69) and (72) follows

$$\bar{J}_{\epsilon_0} \cap \left\{ |\text{enc}(B)| \geq \pi T^2 \right\} \subseteq \left(\bigcup_{i=1}^N J(\epsilon_i) \cap \left\{ |\text{enc}(B)| \geq \pi T^2 \right\} \right) \cup Q_{\frac{1}{CT}, 1}^c.$$

From Lemma 7, the bound on $\mathbb{P}(Q_{f,g})$ given by Lemma 5, and (73), it follows that

$$\begin{aligned} & \mathbb{P}\left(\bar{J}_{\epsilon_0} \cap \left\{ |\text{enc}(B)| \geq \pi T^2 \right\}\right) \\ & \leq CT^2 \log T \exp \left\{ -2\pi^2 T - 2^{-9} T^{\frac{2}{3} + 3\epsilon_0} \right\} \\ & \quad + \left(\frac{32\sqrt{2}}{\sqrt{\pi}} \sqrt{CT}^{\frac{3}{2}} + \frac{16}{\sqrt{\pi}} T^{\frac{1}{2}} \right) \exp \left\{ -\frac{C}{128} T \right\}. \end{aligned}$$

The second term here is negligible provided that $C > 20\pi^2$. Choosing $C_3 = 2C$ yields the statement in the Proposition. $\square$

3 Proof of Theorem 1

We require the following result.

Lemma 8 *Let K denote a planar compact convex set whose area exceeds πT^2 . Then*

$$\text{arcl}(\partial K)^2 > 4\pi^2 T^2 + \pi^2 (\text{R}_{\text{out}}(K) - \text{R}_{\text{in}}(K))^2.$$

Proof The result is implied by Bonnesen's inequality, as it is stated in the Theorem of Subsection 1.3.1, on page 3 of [2]. $\square$

Proof of Theorem 1. We will prove that, for $\epsilon \in (0, 1/6)$ and all sufficiently high T ,

$$\begin{aligned} & \mathbb{P}\left(\left\{ \text{R}_{\text{out}}(B) - \text{R}_{\text{in}}(B) > T^{\frac{2}{3} + \epsilon} \right\} \cap \left\{ |\text{enc}(B)| \geq \pi T^2 \right\}\right) \\ & \leq \exp \left\{ -2\pi^2 T - c_4 T^{\frac{1}{3} + 2\epsilon} \right\}, \end{aligned} \quad (74)$$

for any $c_4 \in (0, \pi^2/32)$. In doing so, we will make use of the polygon P , in the case where $m = \lfloor T^{\frac{1}{3}} \rfloor$ and $t' = 0$.

Definition 2 *Let $\epsilon_0, \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4$ be positive constants that satisfy the following bounds: $\epsilon_0 < 1/3$, $2\epsilon_2 > \epsilon_1$, $\epsilon_1 < 1/6$, $\epsilon_2 < 1/3$, $\epsilon_3 > \epsilon_2/4$, $\epsilon_4 > 2\epsilon_0$.*

- Let H_1 denote the event that

$$66\pi^3 T^2 \geq |\text{conv}(P)| \geq \pi T^2 - T^{\frac{4}{3} + \epsilon_1}.$$

- Let H_2 be given by

$$H_2 = \left\{ R_{\text{out}}(\text{conv}(P)) - R_{\text{in}}(\text{conv}(P)) \leq T^{\frac{2}{3} + \epsilon_2} \right\}.$$

- Let H_3 be given by

$$H_3 = \left\{ \sup_{k \in \{0, \dots, m-1\}} d\left(\partial(\text{conv}P), B\left(\frac{kT}{m}\right)\right) \leq T^{\frac{2}{3} + \epsilon_3} \right\}.$$

- Let H_4 denote the event that, for each $i \in \{0, \dots, m-1\}$,

$$\sup_{t \in [0, T/m]} \left| B\left(\frac{(i-1)T}{m} + t\right) - B\left(\frac{(i-1)T}{m}\right) \right| \leq T^{\frac{2}{3} + \epsilon_4}.$$

Note that

$$\begin{aligned} & \mathbb{P}\left(H_1^c \cap \left\{ |\text{enc}(B)| \geq \pi T^2 \right\}\right) \\ & \leq \mathbb{P}\left(\left\{ |\text{conv}(P)| < \pi T^2 - T^{\frac{4}{3} + \epsilon_1} \right\} \cap \left\{ |\text{enc}(B)| \geq \pi T^2 \right\}\right) \\ & \quad + \mathbb{P}\left(|\text{conv}(P)| > 66\pi^3 T^2\right). \end{aligned} \quad (75)$$

The first event after the inequality in (75) coincides with $H(\epsilon_1)^c \cap \{|\text{enc}(B)| \geq \pi T^2\}$ as it appears in Lemma 4, with t' set equal to zero. Applying this lemma with the choice $\epsilon = 0$, we deduce that, for sufficiently high values of T , the first term on the right-hand-side of (75) is bounded above by

$$\exp \left\{ -2\pi^2 T - c_1 T^{\frac{1}{3} + 2\epsilon_1} \right\}. \quad (76)$$

Note also that, if $|\text{conv}(P)| > 66\pi^3 T^2$, then

$$\sup_{s, t \in [0, T]} |B(s) - B(t)|^2 \geq \text{diam}(\text{conv}(P))^2 \geq \frac{4}{\pi} |\text{conv}(P)| \geq 4 \cdot 66\pi^2 T^2, \quad (77)$$

the first inequality being valid because the endpoints of the longest diameter of $\text{conv}(P)$ are vertices $B(t_1)$ and $B(t_2)$ of P , the second being the standard isoperimetric inequality. Note that (77) implies the occurrence of $Q_{T, T\pi\sqrt{8.33}}^c$. Thus, we learn from Lemma 5 that

$$\mathbb{P}\left(|\text{conv}(P)| > 6\pi^3 T^2\right) \leq \mathbb{P}\left(Q_{T, T\pi\sqrt{8.33}}^c\right) \leq C \exp \left\{ -\frac{33}{16} \pi^2 T \right\}, \quad (78)$$

By (75), the bound (76) on the first term on its right-hand-side, and (78), we have that, for high values of T ,

$$\mathbb{P}\left(H_1^c \cap \left\{ |\text{enc}(B)| \geq \pi T^2 \right\}\right) \leq \exp \left\{ -2\pi^2 T - \frac{c_1}{2} T^{\frac{1}{3} + 2\epsilon_1} \right\}. \quad (79)$$

We assert that

$$\mathbb{P}\left(H_1 \cap H_2^c\right) \leq \exp\left\{-2\pi^2 T - c_2 T^{\frac{1}{3}+2\epsilon_2}\right\}, \quad (80)$$

for any choice of $c_2 \in (0, \pi^2/2)$. To show this, we choose δ to satisfy $\delta < 2/3$ and $\delta \in (\epsilon_1, 2\epsilon_2)$. Note firstly that, since $\delta \in (\epsilon_1, 2/3)$,

$$\pi T^2 - T^{\frac{4}{3}+\epsilon_1} \geq \pi(T - T^{\frac{1}{3}+\delta})^2,$$

for T sufficiently high. The occurrence of H_1 therefore implies that

$$|\text{conv}(P)| \geq \pi(T - T^{\frac{1}{3}+\delta})^2, \quad (81)$$

whereas, on the event H_2^c ,

$$R_{\text{out}}(\text{conv}(P)) - R_{\text{in}}(\text{conv}(P)) > T^{\frac{2}{3}+\epsilon_2}.$$

Using the fact that $\delta < 2/3$, we may apply Lemma 8 to find a lower bound on the arclength of the convex hull of the polygon P in this eventuality:

$$\text{arcl}(\text{conv}(P))^2 \geq 4\pi^2 \left(T - T^{\frac{1}{3}+\delta}\right)^2 + \pi^2 T^{\frac{4}{3}+2\epsilon_2}. \quad (82)$$

From (82) follows

$$\begin{aligned} \text{arcl}(\text{conv}(P)) &\geq 2\pi(T - T^{\frac{1}{3}+\delta})\sqrt{1 + \frac{1}{4}T^{\frac{-2}{3}+2\epsilon_2}} \\ &\geq 2\pi\left(T - T^{\frac{1}{3}+\delta}\right)\left(1 + \left(\frac{1}{8} - o(1)\right)T^{\frac{-2}{3}+2\epsilon_2}\right), \end{aligned} \quad (83)$$

where the facts that $\delta < 2/3$ and $\epsilon_2 < 1/3$ were used in successive inequalities. Since $2\epsilon_2 > \delta$, we deduce that

$$\text{arcl}(\text{conv}(P)) \geq 2\pi T + cT^{\frac{1}{3}+2\epsilon_2}, \quad (84)$$

for any $c \in (0, \pi/4)$. Recalling Definition 1, we find that

$$\begin{aligned} \sum_{i=1}^m L_i^2 &\geq \frac{\left(\sum_{i=1}^m L_i\right)^2}{m} \geq \frac{\left(\text{arcl}(\text{conv}(P))\right)^2}{m} \\ &\geq \frac{\left(2\pi T + cT^{\frac{1}{3}+2\epsilon_2}\right)^2}{m} \geq \frac{4\pi^2 T^2 + 4\pi c T^{\frac{4}{3}+2\epsilon_2}}{m}. \end{aligned}$$

Thus,

$$\sum_{i=1}^{2m} E_i^2 \geq 4\pi^2 T + 4\pi c T^{\frac{1}{3}+2\epsilon_2}, \quad (85)$$

where the collection of random variables $\{E_i : i \in \{1, \dots, 2m\}\}$ was introduced in (28). The left-hand-side of (85) having the χ^2 -distribution with $2m - 2$ degrees of freedom, and m being at most $T^{1/3}$, it follows by (36) that the probability of the occurrence of (85) is at most

$$(CT)^{\frac{1}{2}T^{1/3}} \exp \left\{ -2\pi^2 T - 2\pi c T^{1/3+2\epsilon_2} \right\} + \exp \{ -CT/4 \},$$

for any constant C . Since c may be chosen to lie arbitrarily close to $\pi/4$, we have shown that (80) holds.

We will now show that

$$\mathbb{P}(H_1 \cap H_2 \cap H_3^c) \leq \exp \left\{ -2\pi^2 T - c_3 T^{\frac{1}{2}} \right\}, \quad (86)$$

for any $c_3 \in (0, 2\pi)$. We do so by proving the inclusion

$$H_1 \cap H_2 \cap H_3^c \subseteq \left\{ \sum_{i=1}^m L_i - \text{arcl}(\partial(\text{conv}P)) > T^{\frac{1}{2}} \right\}, \quad (87)$$

for which purpose, we require a lemma.

Lemma 9 *Let P denote a planar polygon, with vertex set $\{p_i : i \in \{1, \dots, m\}\}$ (so that its arclength $\text{arcl}(P)$ is given by the sum of its edge-lengths $|p_{i+1} - p_i|$ for $i \in \{1, \dots, m\}$). Then*

$$\text{arcl}(P) \geq \text{arcl}(\partial(\text{conv}P)) + (\sqrt{5} - 2) \min \left\{ \frac{2R^2}{Q}, R \right\},$$

where $R = \sup_{i \in \{1, \dots, m\}} d(p_i, \partial(\text{conv}P))$, and where Q is equal to the supremum of the lengths of line segments in $\partial(\text{conv}P)$.

Proof. Let L denote the collection of line segments that comprise $\partial(\text{conv}P)$. Write $\partial_{\text{ext}}(P)$ for the exterior boundary of the polygon P . That is, $\partial_{\text{ext}}(P) = \partial(\text{enc}(P)^c)$. Note that the set $\text{conv}(P) \setminus \text{enc}(P)$ is comprised of a finite number of connected components. The boundary of each component consists of the union of a line segment $l \in L$, and a polygonal path lying in $\partial_{\text{ext}}(P)$ whose endpoints coincide with those of l . We denote this polygonal path by l_{ext} . In the case that $l \in L$ does not arise from any such component, we set $l_{\text{ext}} = l$. Note that, for any pair $l', \tilde{l} \in L$, the set $l'_{\text{ext}} \cap \tilde{l}_{\text{ext}}$ has at most finitely many elements. Note further that $|l_{\text{ext}}| \geq |l|$ for each $l \in L$, where we use $|\cdot|$ to denote the length of a line segment, or of a finite union of line segments.

Let $k \in \{1, \dots, m\}$ satisfy

$$R = d(p_k, \partial(\text{conv}P)). \quad (88)$$

We distinguish two cases, according to whether or not

$$d(p_k, \partial_{\text{ext}}(P)) \leq \frac{R}{2}. \quad (89)$$

Supposing that (89) holds, let $x \in \partial_{\text{ext}}(P)$ satisfy

$$d(x, p_k) \leq \frac{R}{2}. \quad (90)$$

Let $l^* = [p_i, p_j]$ denote the element of L for which $x \in l^*$. By using (88) and (90), we find that

$$d(x, l^*) > \frac{R}{2}. \quad (91)$$

Note also that

$$|l_{\text{ext}}^*| \geq |x - p_i| + |p_j - x| \geq |q - p_i| + |q - p_j|,$$

where q denotes the point at distance $\frac{R}{2}$ from l^* whose projection onto this line segment is its midpoint. In the latter inequality, we invoked (91). Thus,

$$|l_{\text{ext}}^*| \geq \sqrt{|p_j - p_i|^2 + R^2} \quad (92)$$

$$= |p_j - p_i| \sqrt{1 + \frac{R^2}{|p_j - p_i|^2}}. \quad (93)$$

If

$$|p_j - p_i| > 2R, \quad (94)$$

then

$$|l_{\text{ext}}^*| \geq |p_i - p_j| + 2(\sqrt{5} - 2) \frac{R^2}{|p_j - p_i|} \geq |p_i - p_j| + 2(\sqrt{5} - 2) \frac{R^2}{Q}. \quad (95)$$

In the first inequality of (95), we used (94) and the fact that

$$\sqrt{1+x} \geq 1 + 2(\sqrt{5} - 2)x$$

for $x \in [0, 1/4]$ to bound below the term appearing in (93). In the second inequality of (95), we used the fact that $|l^*| \leq Q$ (which follows from the definition of Q). Thus, provided that (89) and (94) hold, we have that

$$\begin{aligned} \text{arcl}(P) &\geq \sum_{l \in L} |l_{\text{ext}}| \geq \sum_{l \in L \setminus \{l^*\}} |l| + |p_i - p_j| + 2(\sqrt{5} - 2) \frac{R^2}{Q} \\ &= \text{arcl}(\partial(\text{conv} P)) + 2(\sqrt{5} - 2) \frac{R^2}{Q}. \end{aligned} \quad (96)$$

where the second inequality is due to the fact that $|l_{\text{ext}}| \geq |l|$ for each $l \in L$. Note that, if (94) fails, then

$$|l_{\text{ext}}^*| - |p_i - p_j| \geq \sqrt{|p_i - p_j|^2 + R^2} - |p_i - p_j| \geq (\sqrt{5} - 2)R, \quad (97)$$

where the first inequality follows from (92) and where the second is obtained by minimising the value of the term $\sqrt{|p_i - p_j|^2 + R^2} - |p_i - p_j|$ subject to the constraint that (94) does not hold. The inequality arising from (97) may replace (95) in deriving (96). We have established the statement of the lemma in the event that (89) holds. In the other case, there exists a union $\hat{l}$ of line segments in P such that $p_k \in \hat{l}$, $|\hat{l}| \geq \frac{R}{2}$, and $\hat{l} \cap \partial_{\text{ext}}(P) = \emptyset$. Given that $|\partial_{\text{ext}}(P)| \geq \text{arcl}(\partial(\text{conv}P))$ (which follows from $|l_{\text{ext}}| \geq |l|$ for each $l \in L$), we find that

$$\text{arcl}(P) \geq |\partial_{\text{ext}}(P)| + |\hat{l}| \geq \text{arcl}(\partial(\text{conv}P)) + \frac{R}{2}.$$

Given that $\frac{1}{2} > \sqrt{5} - 2$, this establishes the statement of the lemma in the case that (89) fails, thereby completing the proof of the lemma. $\square$

In applying Lemma 9, we firstly find an upper bound on the quantity Q appearing in its statement, for the polygon P under discussion. To be specific, we now show that, provided that $H_1 \cap H_2$ occurs, then, for any line segment L in the convex boundary of P ,

$$|L| \leq C_4 T^{\frac{5}{6} + \frac{\epsilon_2}{2}}, \quad (98)$$

where C_4 is any constant exceeding $2^{3/2}(66)^{1/4}\sqrt{\pi}$. Indeed, if $H_1 \cap H_2$ occurs, there exist two planar circles J_1 and J_2 , satisfying

$$\text{rad}(J_1) \in \left[\frac{T}{2}, \pi\sqrt{66}T\right], \text{rad}(J_2) \in \left[\text{rad}(J_1), \text{rad}(J_1) + T^{\frac{2}{3} + \epsilon_2}\right],$$

and $\text{enc}(J_1) \subseteq \text{enc}(J_2)$, such that $\partial(\text{conv}P)$ lies in $\text{enc}(J_2) \setminus \text{enc}(J_1)$.

The length of the longest line segment L that lies in $\text{enc}(J_2) \setminus \text{enc}(J_1)$ for a pair of circles J_1 and J_2 satisfying these conditions is attained when $\text{rad}(J_1)$ and $\text{rad}(J_2)$ are maximal, with the relative positions of J_1, J_2 and L resembling the picture in Figure 6. Thus, this length is bounded above by

$$\begin{aligned} & 2\sqrt{\left(\pi\sqrt{66}T + T^{\frac{2}{3} + \epsilon_2}\right)^2 - 66\pi^2 T^2} \\ & \leq 2\sqrt{2\pi\sqrt{66}T^{\frac{5}{3} + \epsilon_2} + T^{\frac{4}{3} + 2\epsilon_2}} \leq 2^{3/2}(66)^{1/4}\sqrt{\pi}(1 + o(1))T^{\frac{5}{6} + \frac{\epsilon_2}{2}}, \end{aligned} \quad (99)$$

the second inequality requiring $\epsilon_2 < 1/3$. This establishes (98).

To apply Lemma 9 and derive (87), note that the occurrence of H_3^c implies that $R > T^{\frac{2}{3} + \epsilon_3}$. It follows from (98) that, in the event of $H_1 \cap H_2$,

we have the inequality, $Q \leq C_4 T^{\frac{5}{6} + \frac{\epsilon_2}{2}}$. Making use of the assumption that $\epsilon_3 > \epsilon_2/4$, we indeed obtain (87) from Lemma 9.

Suppose that the quantity δ chosen before (81) satisfies $\delta < 1/6$ (at this point, we require that $\epsilon_1 < 1/6$). Applying the standard isoperimetric inequality and using (81), which holds provided that H_1 occurs, we find that, on H_1 ,

$$\text{arcl}(\partial(\text{conv} P)) \geq 2\pi T - 2\pi T^{\frac{1}{3} + \delta}.$$

From (96), whose validity we have established whether or not (94) holds, we deduce that

$$\sum_{i=1}^m L_i \geq 2\pi T + (1 - o(1))T^{\frac{1}{2}},$$

since $\delta < 1/6$. By a reprise of the argument that follows (84), the probability of this last inequality is at most $\exp\left\{-2\pi^2 T - 2\pi(1 - o(1))T^{\frac{1}{2}}\right\}$. We have now established (86).

Given that $\epsilon_0 < \min\{1/3, \epsilon_4/2\}$, Lemma 6 implies that $\overline{\mathcal{J}}_{\epsilon_0}^c \subseteq H_4$. Let $\psi(T) = T^{\frac{2}{3}}(T^{\epsilon_2} + 2T^{\epsilon_3} + 2T^{\epsilon_4})$. It is straightforward that

$$H_2 \cap H_3 \cap H_4 \subseteq \left\{R_{\text{out}}(B) - R_{\text{in}}(B) \leq \psi(T)\right\}. \quad (100)$$

From (100), it follows that

$$\begin{aligned} & \mathbb{P}\left(\left\{R_{\text{out}}(B) - R_{\text{in}}(B) > \psi(T)\right\} \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\}\right) \\ & \leq \mathbb{P}\left((H_2 \cap H_3 \cap H_4)^c \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\}\right) \\ & \leq \mathbb{P}\left(H_1^c \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\}\right) + \mathbb{P}(H_1 \cap H_2^c) \\ & \quad + \mathbb{P}(H_1 \cap H_2 \cap H_3^c) + \mathbb{P}\left(\overline{\mathcal{J}}_{\epsilon_0} \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\}\right) \\ & \leq \exp\left\{-2\pi^2 T - \frac{c_1}{2}T^{\frac{1}{3} + 2\epsilon_1}\right\} + \exp\left\{-2\pi^2 T - c_2 T^{\frac{1}{3} + 2\epsilon_2}\right\} \\ & \quad + \exp\left\{-2\pi^2 T - c_3 T^{\frac{1}{2}}\right\} + C_3 T^2 \log T \exp\left\{-2\pi^2 T - 2^{-9} T^{\frac{2}{3} + 3\epsilon_0}\right\}, \end{aligned} \quad (101)$$

where, in the second inequality, we used $H_4^c \subseteq \overline{\mathcal{J}}_{\epsilon_0}$. The bounds (79), (80), (86) and that provided by Proposition 2 were used for the successive terms to obtain the third inequality. For $\epsilon \in (0, 1/6)$, we set $\epsilon_1 = \epsilon + \hat{\epsilon}$, $\epsilon_2 = \epsilon_4 = \epsilon$, $\epsilon_3 = \epsilon/4 + \hat{\epsilon}$ and $\epsilon_0 = \epsilon/2 - \hat{\epsilon}$, where $\hat{\epsilon} > 0$ is chosen small enough that the restrictions on the parameters $\{\epsilon_i : i \in \{0, \dots, 4\}\}$ are satisfied. We obtain

$$\begin{aligned} & \mathbb{P}\left(\left\{R_{\text{out}}(B) - R_{\text{in}}(B) > 3T^{\frac{2}{3} + \epsilon}\right\} \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\}\right) \\ & \leq \exp\left\{-2\pi^2 T - (c_2 - o(1))T^{\frac{1}{3} + 2\epsilon}\right\}, \end{aligned}$$

since $\epsilon \in (0, 1/6)$. From this follows (74), the value of c_4 in (74) arising because we may choose c_2 in (80) to be slightly less than $\pi^2/2$. Note that

$$\begin{aligned} & \left\{ R_{\text{in}}(B) < T - T^{\frac{2}{3}+\epsilon} \right\} \cap \left\{ |\text{enc}(B)| \geq \pi T^2 \right\} \\ & \subseteq \left\{ R_{\text{out}}(B) - R_{\text{in}}(B) > T^{\frac{2}{3}+\epsilon} \right\}, \end{aligned} \quad (102)$$

because $R_{\text{out}}(B) \geq T$ if $|\text{enc}(B)| \geq \pi T^2$.

We obtain

$$\begin{aligned} & \mathbb{P}\left(R_{\text{in}}(X) < T - T^{\frac{2}{3}+\epsilon}\right) \\ &= \frac{\mathbb{P}\left(\left\{R_{\text{in}}(B) < T - T^{\frac{2}{3}+\epsilon}\right\} \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\}\right)}{\mathbb{P}\left(|\text{enc}(B)| \geq \pi T^2\right)} \\ &\leq \exp\left\{-c_4 T^{\frac{1}{3}+2\epsilon} + C_1 T^{\frac{1}{3}} \log T\right\}, \end{aligned}$$

the inequality by means of (74), (102) and Lemma 1. We have proved the first part of the theorem.

To derive the second statement, note that

$$\begin{aligned} & \left\{ R_{\text{out}}(B) > T + 2T^{\frac{2}{3}+\epsilon} \right\} \cap \left\{ |\text{enc}(B)| \geq \pi T^2 \right\} \\ & \subseteq \left\{ R_{\text{in}}(B) > T + T^{\frac{2}{3}+\epsilon} \right\} \end{aligned} \quad (103)$$

$$\cup \left\{ R_{\text{out}}(B) - R_{\text{in}}(B) > T^{\frac{2}{3}+\epsilon} \right\} \cap \left\{ |\text{enc}(B)| \geq \pi T^2 \right\}. \quad (104)$$

Noting that the inequality $R_{\text{in}}(B) > T + T^{\frac{2}{3}+\epsilon}$ implies that $|\text{enc}(B)| \geq \pi T^2 + 2\pi T^{\frac{5}{3}+\epsilon}$, we may use Proposition 1 to bound the probability of the event in (103). We obtain that, for any $\alpha \in (0, 1/3)$,

$$\mathbb{P}\left(R_{\text{in}}(B) > T + T^{\frac{2}{3}+\epsilon}\right) \leq \exp\left\{-2\pi^2 T - T^{\frac{1}{3}+\alpha}\right\}, \quad (105)$$

provided that T is chosen to be sufficiently high.

Using (74) to bound the probability of the event in (104), as well as (105) we find that, for $\epsilon \in (0, 1/6)$ and for all sufficiently high values of T ,

$$\begin{aligned} & \mathbb{P}\left(\left\{R_{\text{out}}(B) > T + 2T^{\frac{2}{3}+\epsilon}\right\} \cap \left\{|\text{enc}(B)| \geq \pi T^2\right\}\right) \\ & \leq \exp\left\{-2\pi^2 T - \frac{c_4}{2} T^{\frac{1}{3}+2\epsilon}\right\}. \end{aligned} \quad (106)$$

From (106) and Lemma 1, we deduce the second statement of the theorem.

□




Figure 6: Computing the length $|L|$ in (109)

Lemma 10 *Let K denote a planar compact set. In the case that $R_{\text{in}}(K) > \frac{1}{2}R_{\text{out}}(K)$, we have that*

$$\mathcal{L}(\text{conv}K) \leq 4\sqrt{R_{\text{in}}(K)(R_{\text{out}}(K) - R_{\text{in}}(K))}.$$

Proof Given that K is compact, we may locate circles J_1 and J_2 satisfying

$$\begin{aligned} \text{enc}(J_1) \subseteq K \subseteq \text{enc}(J_2), \quad \text{rad}(J_2) &= R_{\text{out}}(K) \\ \text{and } \text{rad}(J_1) &= R_{\text{in}}(K). \end{aligned} \quad (107)$$

Note that any line segment L lying in $\partial(\text{conv}K)$ satisfies

$$L \subseteq \text{int}(\text{enc}(J_1))^c \cap \text{enc}(J_2), \quad (108)$$

where $\text{int}(A)$ denotes the interior of the set A . Indeed, if $\text{int}(\text{enc}(J_1)) \cap L \neq \emptyset$, there exists a point lying in $\text{enc}(J_1)$ and in the half-plane whose boundary contains L and is disjoint from $\text{conv}(K)$. This point does not lie in K , implying that $\text{enc}(J_1) \not\subseteq K$. The endpoints of L lie in $\text{enc}(J_2)$, and thus, so does L , by the convexity of $\text{enc}(J_2)$.

Given that $R_{\text{in}}(K) > \frac{1}{2}R_{\text{out}}(K)$, the supremum of the lengths of line segments L satisfying (108) over the set of pairs of circles (J_1, J_2) satisfying (107) is achieved when J_1 and J_2 touch at a point, with L being the line segment tangent to J_1 at the diametrically opposed point and having endpoints in J_2 (see Figure 6). For this choice of line segment, $|L|$ satisfies

$$|L| = 2\sqrt{R_{\text{out}}(K)^2 - \left(2R_{\text{in}}(K) - R_{\text{out}}(K)\right)^2} \quad (109)$$

$$= 4\sqrt{R_{\text{in}}(K)(R_{\text{out}}(K) - R_{\text{in}}(K))},$$

as required. $\square$

Proof of Corollary 1 From the first part of Lemma 10 follows the inclusion

$$\begin{aligned} & \left\{ \mathcal{L}(\text{conv} B) > T^{\frac{5}{6}+\epsilon} \right\} \\ \subseteq & \left\{ R_{\text{in}}(B) > C_5 T \right\} \cup \left\{ R_{\text{out}}(B) - R_{\text{in}}(B) \geq \frac{1}{16C_5} T^{\frac{2}{3}+2\epsilon} \right\}, \end{aligned} \quad (110)$$

for any fixed $C_5 > 0$. Note that the event $Q_{f,g}^c$ occurs when the choices $f(T) = T$ and $g(T) = R_{\text{in}}(B)$ are made. From Lemma 5, it follows that

$$\mathbb{P}(R_{\text{in}}(B) > C_5 T) \leq \exp \left\{ -\frac{C_5^2}{128} T \right\}. \quad (111)$$

We deduce that

$$\begin{aligned} & \mathbb{P} \left(\left\{ \mathcal{L}(\text{conv} B) > T^{\frac{5}{6}+\epsilon} \right\} \cap \left\{ |\text{enc}(B)| \geq \pi T^2 \right\} \right) \\ \leq & \mathbb{P} \left(\left\{ R_{\text{out}}(B) - R_{\text{in}}(B) > \frac{1}{16C_5} T^{\frac{2}{3}+2\epsilon} \right\} \cap \left\{ |\text{enc}(B)| \geq \pi T^2 \right\} \right) \\ & + \mathbb{P}(R_{\text{in}}(B) > C_5 T) \\ \leq & \exp \left\{ -2\pi^2 T - c_5 T^{\frac{1}{3}+4\epsilon} \right\} + C \exp \left\{ -\frac{C_5^2}{128} T \right\}, \end{aligned} \quad (112)$$

where c_5 is a constant satisfying $c_5 \leq c_4/(16C_5)^2$, and where $\epsilon \in (0, 1/12)$. (Recall that the constant c_4 appeared in (74).) The first inequality is a consequence of (110). In the second, the bounds (111) and (74) were used. The choice $C_5 > 16\pi$ ensures that the second term in (112) is negligible. Thus, c_5 may be chosen to be any value satisfying $c_5 \leq 1/(32(16)^4) = 2^{-21}$. The first part of the corollary follows by applying Lemma 1.

For the second part of the corollary, note that, if $R_{\text{out}}(B) \leq T + T^{\frac{2}{3}+\epsilon}$, then $\text{arcl}(\partial(\text{conv} B)) \leq 2\pi(T + T^{\frac{2}{3}+\epsilon})$, because the arclength of the boundary of a planar convex set is monotone under containment [6]. Thus, the second part of Theorem 1 yields the result. $\square$

Acknowledgments: We are grateful to Senya Shlosman for proposing the model to us, and thank him in particular for presenting heuristic arguments about the nature of its deviation. We thank Manjunath Krishnapur, Ben Hough and Dapeng Zhan for helpful comments on a draft version of the paper.

References

- [1] Kenneth S. Alexander. Cube-root boundary fluctuations for droplets in random cluster models. *Comm. Math. Phys.*, 224(3):733–781, 2001.

- [2] Yu. D. Burago and V. A. Zalgaller. *Geometric inequalities*, volume 285 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1988. Translated from the Russian by A. B. Sosinskiĭ, Springer Series in Soviet Mathematics.
- [3] Amir Dembo and Ofer Zeitouni. Large deviations and applications. In *Handbook of stochastic analysis and applications*, volume 163 of *Statist. Textbooks Monogr.*, pages 361–416. Dekker, New York, 2002.
- [4] Richard Durrett. *Probability: theory and examples*. Duxbury Press, Belmont, CA, second edition, 1996.
- [5] Ioannis Karatzas and Steven E. Shreve. *Brownian motion and stochastic calculus*, volume 113 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1988.
- [6] L. A. Santaló. Integral geometry. In *Studies in Global Geometry and Analysis*, pages 147–193. Math. Assoc. Amer. (distributed by Prentice-Hall, Englewood Cliffs, N.J.), 1967.
- [7] Hasan B. Uzun and Kenneth S. Alexander. Lower bounds for boundary roughness for droplets in Bernoulli percolation. *Probab. Theory Related Fields*, 127(1):62–88, 2003.
- [8] S. R. S. Varadhan. *Large deviations and applications*, volume 46 of *CBMS-NSF Regional Conference Series in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1984.
- [9] David Williams. *Probability with martingales*. Cambridge Mathematical Textbooks. Cambridge University Press, Cambridge, 1991.
- [10] G. Wulff. Zur Frage der Geschwindigkeit des Wachstums und der Auflösung der Krystallflächen. *Z. Kryst.*, 34:449–530, 1901.